

Chapter 3. Curving

1	
2	3.1 The Schwarzschild Metric 3-1
3	3.2 Mass in Units of Length 3-7
4	3.3 The Global Schwarzschild r -Coordinate 3-11
5	3.4 The Global Schwarzschild t -Coordinate 3-17
6	3.5 Constructing the Global Schwarzschild Map of
7	Events 3-18
8	3.6 The Spacetime Slice 3-22
9	3.7 Light Cone Diagram on an $[r, t]$ Slice 3-24
10	3.8 Inside the Event Horizon: A Light Cone Diagram on an
11	$[r, \phi]$ Slice 3-27
12	3.9 Outside the Event Horizon: An Embedding Diagram on
13	an $[r, \phi]$ Slice 3-29
14	3.10 Room and Worldtube 3-33
15	3.11 Exercises 3-35
16	3.12 References 3-41

- 17 • *General relativity describes only tiny effects, right?*
- 18 • *What does “curvature of spacetime” mean?*
- 19 • *What tools can I use to visualize spacetime curvature?*
- 20 • *Do people at different r -coordinates near a black hole age differently?*
- 21 • *If so, can they feel the slowing down/speeding up of their aging?*
- 22 • *What is the “event horizon,” and what weird things happen there?*
- 23 • *Do funnel diagrams describe the gravity field of a black hole?*

CHAPTER

3

25

Curving

Edmund Bertschinger & Edwin F. Taylor *

26 *In my talk ... I remarked that one couldn't keep saying*
 27 *"gravitationally completely collapsed object" over and over.*
 28 *One needed a shorter descriptive phrase. "How about black*
 29 *hole?" asked someone in the audience. I had been searching*
 30 *for just the right term for months, mulling it over in bed, in*
 31 *the bathtub, in my car, wherever I had quiet moments.*
 32 *Suddenly this name seemed exactly right. ... I decided to be*
 33 *casual about the term "black hole," dropping it into [a later]*
 34 *lecture and the written version as if it were an old family*
 35 *friend. Would it catch on? Indeed it did. By now every*
 36 *schoolchild has heard the term.*

37

—John Archibald Wheeler with Kenneth Ford

3.1 ■ THE SCHWARZSCHILD METRIC

39 *Spherically symmetric massive center of attraction?*
 40 *The Schwarzschild metric describes the curved, empty spacetime around it.*

Einstein to
 Schwarzschild:
 "splendid."

41 In late 1915, within a month of the publication of Einstein's general theory of
 42 relativity and just a few months before his own death from a battle-related
 43 illness, Karl Schwarzschild (1873-1916) derived from Einstein's field equations
 44 the metric for spacetime surrounding the spherically symmetric black hole.
 45 Einstein wrote to him, "I had not expected that the exact solution to the
 46 problem could be formulated. Your analytic treatment of the problem appears
 47 to me splendid."

Orbits stay in a plane.

48 An isolated satellite zooms around a spherically symmetric massive body.
 49 After a few orbits we discover that the satellite's motion stays confined to the
 50 initial plane determined by the satellite's position, its direction of motion, and
 51 the center of the attracting body. Why? The reason is simple: symmetry! With

*Draft of Second Edition of *Exploring Black Holes: Introduction to General Relativity*
 Copyright © 2017 Edmund Bertschinger, Edwin F. Taylor, & John Archibald Wheeler. All
 rights reserved. This draft may be duplicated for personal and class use.

3-2 Chapter 3 Curving

Box 1. Metric in Polar Coordinates for Flat Spacetime




FIGURE 1 Spatial separation between two points in polar coordinates.

The metric for flat spacetime is:

$$d\tau^2 = dt^2 - ds^2 \quad (\text{flat spacetime}) \quad (1)$$

where ds is the spatial separation, expressed in Cartesian coordinates as

$$ds^2 = dx^2 + dy^2 \quad (\text{flat space}) \quad (2)$$

We look for a similar ds expression for two adjacent events numbered 1 and 2, events separated by polar coordinate increments dr and $d\phi$ (Figure 1).

Draw little arcs of different radii through events 1 and 2 to form a tiny box, shown in the magnified inset. The squared spatial

separation between events 1 and 2 is—approximately—the sum of the squares of two adjacent sides of the little box. For a differential $d\phi$, we are entitled to express the differential space separation between event 1 and event 2 by the formula

$$ds^2 = dr^2 + r^2 d\phi^2 \quad (\text{flat space}) \quad (3)$$

This squared spatial separation is the space part of the squared wristwatch time differential for flat spacetime

$$d\tau^2 = dt^2 - dr^2 - r^2 d\phi^2 \quad (\text{flat spacetime}) \quad (4)$$

This derivation is valid only when $d\phi$ is small—vanishingly small in the calculus sense—so that the differential segment of arc $r d\phi$ is indistinguishable from a straight line. There is no such limitation to differentials for rectangular Cartesian space coordinates in flat spacetime, so each d for differential in (2) can be expanded to Δ , as it was in Section 1.10.

From Einstein’s general relativity equations, Schwarzschild derived a generalization of (4) that goes beyond flat spacetime and describes curved spacetime in the vicinity of a spherically symmetric (thus non-spinning) uncharged black hole.

52 respect to this initial plane there is no distinction between “up out of” and
 53 “down out of” the plane, so the satellite cannot choose either and must remain
 54 in that plane. The limitation of isolated particle and light motion to a single
 55 plane greatly simplifies our analysis of physical events in this book.

56 We use polar coordinates (r, ϕ) for the black hole (Box 1), because polar
 57 coordinates reflect its symmetry on a plane through the black hole’s center;
 58 Cartesian coordinates (x, y) do not.

59 Think of two adjacent events that lie on our equatorial r, ϕ plane through
 60 the center of the black hole. These events have differential coordinate
 61 separations dt , dr , and $d\phi$. The **Schwarzschild metric** gives us the invariant
 62 $d\tau$ between this pair of events:

Schwarzschild
timelike metric

$$d\tau^2 = \left(1 - \frac{2M}{r}\right) dt^2 - \frac{dr^2}{\left(1 - \frac{2M}{r}\right)} - r^2 d\phi^2 \quad (\text{timelike}) \quad (5)$$

$$-\infty < t < \infty \quad \text{and} \quad 0 < r < \infty \quad \text{and} \quad 0 \leq \phi < 2\pi$$

64 Equation (5) is the *timelike* form of the Schwarzschild metric, whose left side
 65 gives us the invariant *differential wristwatch time* $d\tau$ of a free stone that moves
 66 between a pair of adjacent events for which the magnitude of the first term on

Section 3.1 The Schwarzschild Metric **3-3**

Schwarzschild
spacelike metric

67 the right side is greater than the magnitude of the last two terms. In contrast,
68 think of a pair of events for which the magnitude of the last two terms on the
69 right predominate. Then the invariant *differential ruler distance* $d\sigma$ between
70 these events is given by the *spacelike* form of the Schwarzschild metric:

$$d\sigma^2 = -d\tau^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \frac{dr^2}{\left(1 - \frac{2M}{r}\right)} + r^2 d\phi^2 \quad (\text{spacelike}) \quad (6)$$

$$-\infty < t < \infty \quad \text{and} \quad 0 < r < \infty \quad \text{and} \quad 0 \leq \phi < 2\pi$$

71
72 Neither a stone nor a light flash can move between an adjacent pair of events
73 with spacelike separation. Instead, the separation $d\sigma$ represents a differential
74 ruler distance between two events. To make use of global metrics (5) and (6),
75 we need to define carefully the meaning of global coordinates t , r , and ϕ .
76 Section 3.2 shows how to measure mass in meters, so that $2M/r$ becomes
77 unitless, as it must in order to subtract it from the unitless number one in the
78 expression $(1 - 2M/r)$.

Meaning of
"global metric"

79 **Comment 1. Terminology: global metric**
80 We refer to either expression (5) or (6) as a *global metric*. Professional general
81 relativists call these expressions *line elements*; they reserve the term *metric* for
82 the collection of coefficients of the differentials—such as $(1 - 2M/r)$, the
83 coefficient of dt^2 . We find the term *metric* to be simple, short, and clear; so in
84 this book we use a slightly-deviant terminology and call an expression like (5) or
85 (6) the **global metric**.

Definition: **invariant**
in general relativity

86 **DEFINITION 1. Invariant (general relativity)**
87 Section 1.2 defined an *invariant* in special relativity as a quantity that
88 has the same value when calculated using different *local inertial*
89 coordinates. An **invariant** in general relativity is a quantity that has the
90 same value when calculated using different *global* coordinate systems.
91 Equations (5) and (6) calculate invariants $d\tau$ and $d\sigma$, respectively, using
92 Schwarzschild global coordinates. Box 3 in Section 7.5 shows that an
93 infinite number of global coordinate systems exist for the non-spinning
94 black hole (indeed, for any isolated black hole). Calculation of $d\tau$ using
95 any of these global coordinate systems delivers the same—the
96 invariant!—value of $d\tau$ given by metrics (5) and (6).

Event horizon

97 Two coefficients in the Schwarzschild metric contain the expression
98 $(1 - 2M/r)$, which goes to zero when $r \rightarrow 2M$, thus sending the first metric
99 coefficient to zero on the right side of the metric and the magnitude of the
100 second coefficient to infinity. This warns us about trouble at $r = 2M$, which we
101 describe below. To the global spacetime surface at $r = 2M$ we assign the name
102 **event horizon**, for reasons that will become clear in later sections.

103 It is important to realize how rare and wonderful is the Schwarzschild
104 metric. Einstein's set of field equations is nonlinear and can be solved in

3-4 Chapter 3 Curving

Simple global metrics are rare.

105 simple form only for physical systems with considerable symmetry.
 106 Schwarzschild used the symmetry of an isolated spherical non-spinning center
 107 of attraction in the derivation of his metric. This symmetry is broken—and no
 108 simple global metric exists—when we place a black hole on every street corner,
 109 although in principle a computer can provide a numerical solution of Einstein’s
 110 field equations for any distribution of mass/energy/pressure. It is a measure of
 111 the scarcity of physical systems with simple metrics that almost fifty years
 112 passed before Roy Kerr found a (relatively!) simple metric for a spinning black
 113 hole in 1963 (Chapters 17 through 21).

Schwarzschild description of spacetime is complete.

114 Further investigation shows that the Schwarzschild metric plus the
 115 connectedness (“topology”) of the region provides a *complete* description of
 116 spacetime external to any isolated spherically symmetric, uncharged massive
 117 body—and everywhere around such a black hole except at its central
 118 singularity (at $r = 0$), where spacetime curvature becomes infinite and general
 119 relativity fails. *Every* feature of spacetime around this kind of black hole is
 120 described or implied by the Schwarzschild metric. This one expression tells it
 121 all!

QUERY 1. Flat spacetime as $r \rightarrow \infty$

Show that as $r \rightarrow \infty$ Schwarzschild metric (5) becomes metric (4) for flat spacetime.

Ways in which the Schwarzschild metric makes sense:

126 We will derive the Schwarzschild metric in Chapter 22. Even now,
 127 however, we should not accept it uncritically. Here we check three ways in
 128 which it makes sense.

1. Depends only on r coordinate.

129 **First**, the expression $(1 - 2M/r)$ that appears in both the dt term and
 130 the dr term depends only on the r coordinate, not on the angle ϕ . How come?
 131 Because we are dealing with a spherically symmetric body, an object for which
 132 there is no way to tell one side from the other side or the top from the bottom.
 133 This impossibility is reflected in the absence of any direction-dependent
 134 coefficient in the metric.

2. Goes to inertial metric for zero M .

135 **Second**, the Schwarzschild metric uses coordinates that clearly show
 136 spacetime is flat when $M \rightarrow 0$, that is when there is *no* center of attraction. In
 137 this limit, the Schwarzschild metric (5) goes smoothly into the inertial metric
 138 (4) for flat spacetime.

3. Goes to local inertial metric for large r .

139 **Third**, even when M is nonzero the Schwarzschild metric (5) reduces to a
 140 local flat spacetime metric (4) very far from the black hole. The expression
 141 $(1 - 2M/r) \rightarrow 1$ when $r \rightarrow \infty$.

Schwarzschild metric applies only outside the surface.

142 Timelike and spacelike Schwarzschild metrics (5) and (6) describe the
 143 spacetime *external to any* isolated spherically symmetric, uncharged massive
 144 body. They apply with high precision to spacetime *outside* a slowly revolving
 145 massive object such as Earth or an ordinary star like our Sun. Think of a
 146 stone moving outside such an object; it makes no difference what the
 147 coordinates are inside the attracting spherical body because the stone never
 148 gets there; before it can, it collides with the surface—in the short term, our

Box 2. More About the Black Hole

John Archibald Wheeler adopted the term “black hole” in 1967 (initial quote), but the concept itself is old. As early as 1783, John Michell argued that light must “be attracted in the same manner as all other bodies” and therefore, if the attracting center is sufficiently massive and sufficiently compact, “all light emitted from such a body would be made to return toward it.” Pierre-Simon Laplace came to the same conclusion independently in 1795 and went on to reason that “it is therefore possible that the greatest luminous bodies in the universe are on this very account invisible.”

Michell and Laplace used Isaac Newton’s “action-at-a-distance” theory of gravity in analyzing the escape of light from, or its capture by, an already-existing compact object. But is such a static compact object possible? In 1939, J. Robert Oppenheimer and Hartland Snyder published the first detailed treatment of gravitational collapse within the framework of Einstein’s theory of gravitation. Their paper predicts the central features of a non-spinning black hole.

Ongoing theoretical study has shown that the black hole is the result of natural physical processes. A nonsymmetric collapsing system is not necessarily blown apart by its instabilities but can quickly—in a few seconds measured on a remote clock!—radiate away its turbulence as gravitational waves and settle down into a stable structure.

An uncharged spherically symmetric black hole is completely described by the Schwarzschild metric (plus the spacetime topology), which was derived from Einstein’s field equations by Karl Schwarzschild and published in 1916. The energy of such a non-spinning black hole cannot be milked for use outside its event horizon. For this reason, a non-spinning black hole deserves the name “dead black hole.”

In contrast to the non-spinning dead black hole, the typical black hole, like the typical star, has a spin, sometimes a large

spin. The energy stored in this spin, moreover, is available for doing work: for driving jets of matter and for propelling a spaceship. In consequence, the spinning black hole deserves and receives the name “live black hole.”

The spinning black hole—or any spinning mass—drags everything in its vicinity around with it, including spacetime (Chapters 17 through 21). Near Earth this dragging is a small effect. Theory predicts that, near a rapidly-spinning black hole, such effects can be large, even irresistible, dragging along nearby spaceships no matter how powerful their rockets.

Black holes appear to be divided roughly into two groups, depending on their source: Those that result from the collapse of a single star have several times the mass our Sun. Others formed near the centers of galaxies can be monsters with millions—even billions—of times the mass of our Sun. These black holes may even shape the evolution of galaxies.

In 1963 Roy P. Kerr derived a metric for an uncharged spinning black hole. In 1967 Robert H. Boyer and Richard W. Lindquist devised a simple and convenient global coordinate system for the spinning black hole. In 2000 Chris Doran published the global coordinate system for a spinning black hole that we use in this book. In 1965 Ezra Theodore Newman and others solved the Einstein equations for the spacetime geometry around an *electrically charged* spinning black hole.

Subsequent research shows that for a steady-state black hole of specified mass, charge, and angular momentum, Kerr-Newman geometry is the *most general* solution to Einstein’s field equations. The variety, detail, and beauty of everything that forms or falls into a black hole disappears—at least according to classical (non-quantum) physics—leaving only mass, charge, and angular momentum. John Wheeler summarized this finding in the phrase, “The black hole has no hair,” which is known as the **no-hair theorem**.

149 Sun can be thought of as in equilibrium. The more compact the massive body,
150 however, the larger the external region the stone can explore. Our Sun’s
151 surface is 696 000 kilometers from its center. A cool white dwarf with the mass
152 of our Sun has a surface r -coordinate of about 5000 kilometers, roughly that of
153 Earth. The Schwarzschild metric describes spacetime geometry in the region
154 external to that r -coordinate. A neutron star with the mass of our Sun has a
155 surface r -coordinate of about 10 kilometers—the size of a typical city—so the
156 stone can come even closer and still be “outside,” that is, in the region
157 described correctly by the Schwarzschild metric (if the neutron star is not
158 spinning too fast).

3-6 Chapter 3 Curving

Box 3. Singularities: Fictitious or Real?




FIGURE 2 Polar coordinates on a flat Euclidean surface have a *coordinate singularity* at the center. Obviously $r = 0$ there, but what is its value of ϕ ? That singularity, however, is fictitious because there is no *space* singularity at that point.

How do we know that the blow-up of the term $dr^2/(1 - 2M/r)$ at $r = 2M$ in the Schwarzschild metric does *not* signal a physical singularity? Why is this blow-up no threat to an observer falling through the event horizon—other than its one-way nature and the gradually-increasing tidal forces she feels as she descends? Einstein and others initially thought that the Schwarzschild coordinate singularity at the event horizon had a physical reality, but it does not.

Similarly, how do we know that the blow-up of the term $(1 - 2M/r)dt^2$ at $r = 0$ is lethal to all comers? How can we understand the difference between the two discontinuities in Schwarzschild coordinates?

Draw an analogy to the polar coordinate system (r, ϕ) on a flat Euclidean surface (Figure 2). The radial coordinate of

the origin is clearly $r = 0$, but what is the polar angle ϕ there? *Answer:* The origin is singular in angle ϕ . *Proof:* Start at the right on the horizontal axis with label $\phi = 0$; move leftward along this axis and through the origin at $r = 0$. At this origin the axis label suddenly flips to $\phi = 180^\circ$. There is a discontinuity of ϕ at the origin. The *coordinate* ϕ violates the requirements of uniqueness and smoothness.

The problem here is *not* Euclidean space, it is our silly (r, ϕ) coordinate system. In contrast, Cartesian coordinates $x = r \cos \phi$ and $y = r \sin \phi$ are perfectly unique and continuous at all points on the flat surface, including the origin.

Is there some way to show that there is no physical singularity at the event horizon of a non-spinning black hole? Yes, by finding a coordinate system which is perfectly smooth at the event horizon, in the same way that Cartesian coordinates in Euclidean space are perfectly smooth at the origin. In Chapter 7 we develop what we call *global rain coordinates*. At the event horizon no term blows up in the metric expressed in global rain coordinates. Global rain coordinates assign unique labels to each event and are smooth and continuous at the event horizon and all the way down to (but not including) $r = 0$.

What about the location at the center of a black hole? No coordinate system can be smooth at $r = 0$, because the so-called **Riemann curvature** is infinite there. The Riemann curvature, discovered in the 1860s by mathematician Bernhard Riemann, has a value at every spacetime event that is independent of the coordinate system. The Riemann curvature is infinite only at a physical cusp or singularity, such as the black hole singularity at $r = 0$. In contrast, the Riemann curvature is finite at $r = 2M$.

Schwarzschild describes **all** spacetime around the black hole outside the singularity.

159 A wonderful thing about a black hole is that it has no physical surface and
160 no matter with which to collide. A stone can explore *all* of spacetime (except
161 at $r = 0$) without bumping into a surface—since there is no surface at all.



162 **Objection 1.** How can a black hole have “no matter with which to collide”?
163 If it isn’t made of matter, what is it made of? What happened to the star or
164 group of stars that collapsed to form the black hole? Basically, how can
165 something have mass without being made of matter?



166 We think that everything that collapses into the black hole is effectively still
167 there in some form, inducing the curvature of surrounding spacetime. This
168 mass is crushed into a singularity at the center—along with the probe we

Section 3.2 Mass in Units of Length 3-7

169 sent in to explore it. How do we know this? We don't. What can "crushed to
170 a singularity" possibly *mean*? We don't know. Startling? Crazy? Absurd?
171 Welcome to general relativity!

?

172 **Objection 2.** *The global metric comes from Einstein's equations, which*
173 *you say we will derive in Chapter 22. In the meantime you give us only*
174 *global metrics. Why should we believe you, and why are you keeping the*
175 *fundamental equations from us?*

!

176 Einstein's equations are most economically expressed in advanced
177 mathematics such as *tensors*, and deriving a global metric from them is a
178 bit tricky. In contrast, the global metric expresses itself in differentials, the
179 working mathematics of most technical professions, and leads directly to
180 measurable quantities: wristwatch time and ruler distance. We choose to
181 start with the directly useful.

182 Next we examine the meaning of mass in units of length, so that the
183 expression $1 - 2M/r$ in both the first and second term in the metric
184 coefficients can have the same units, namely no units at all.

3.2 ■ MASS IN UNITS OF LENGTH

186 *Want to reduce clutter in the metric? Then measure mass in meters!*

187 The description of spacetime near any gravitating body is simplest when we
188 express the mass M of that body in spatial units—in meters or kilometers.
189 This section derives the conversion factor between, for example, kilograms and
190 meters.

Measure mass
in meters.

191 Earlier we wanted to measure space and time in the same unit (Section
192 1.2), so we used the conversion factor c , the speed of light. Conversion from
193 kilograms to meters is not so simple. Nevertheless, here too Nature provides a
194 conversion factor, a combination of the speed of light and Newton's **universal**
195 **gravitation constant** G .

196 Newton's theory of gravitation predicts that the gravitational force
197 between two spherically symmetric masses M_{kg} and m_{kg} is proportional to the
198 product of these masses and inversely proportional to the square of the
199 Euclidean distance r between their centers:

$$F_{\text{Newtons}} = -\frac{GM_{\text{kg}}m_{\text{kg}}}{r^2} \quad (\text{Newton, conventional units}) \quad (7)$$

200 In this equation G is the "constant of proportionality," whose units depend on
201 the units with which mass and spatial separation are measured. The numerical
202 value of G in conventional units is:

$$G = 6.67 \times 10^{-11} \frac{\text{meter}^3}{\text{kilogram second}^2} \quad (8)$$

3-8 Chapter 3 Curving

Numerical
values of G

203 Divide G by the square of the speed of light c^2 to find the conversion factor
204 that translates the conventional unit of mass, the kilogram, into what we have
205 already chosen to be the natural unit, the meter:

$$\begin{aligned} \frac{G}{c^2} &= \frac{6.67 \times 10^{-11} \frac{\text{meter}^3}{\text{kilogram second}^2}}{8.9876 \times 10^{16} \frac{\text{meter}^2}{\text{second}^2}} \\ &= 7.42 \times 10^{-28} \frac{\text{meter}}{\text{kilogram}} \end{aligned} \quad (9)$$

206 Now convert from mass M_{kg} measured in conventional units of kilograms to
207 mass M in meters by multiplication with this conversion factor:

$$M \equiv \frac{G}{c^2} M_{\text{kg}} = \left(7.42 \times 10^{-28} \frac{\text{meter}}{\text{kilogram}} \right) M_{\text{kg}} \quad (10)$$

Mass in meters
unclutters equations.

208 *Why* make this conversion? Because it allows us to get rid of the symbols G
209 and c^2 that otherwise clutter up our equations.

210 Table 1 displays in both kilograms and meters the masses of Earth, Sun,
211 the huge spinning black hole at the center of our galaxy, and the mass of an
212 even larger black hole in a nearby galaxy. For each of these objects the global
213 r -coordinate of the event horizon is twice its mass in meters. To express their
214 masses in meters cuts planets and stars down to size!

?

215 **Objection 3.** *This is nuts! Stars and planets are not the same as space.*
216 *No twisting or turning on your part can make mass and distance the same.*
217 *How can you possibly propose to measure mass in units of meters?*

!

218 True, mass is not the same as spatial separation. Neither is time the same
219 as space: The separation between clock ticks is different from meterstick
220 lengths! Nevertheless, we have learned to use the conversion factor c to
221 measure both time and space in the same unit: light-years of spatial
222 separation and years of time, for example, or meters of spatial coordinate
223 separation and meters of light-travel time. Payoff? The result simplifies our
224 equations.

225 There are two primary birthplaces for black holes: The first is the collapse
226 of a single star, which produces a black hole with mass equal to a modest
227 multiple of the mass of our Sun. The second birthplace is accumulation in a
228 galaxy, which produces a black hole with mass equal thousands to billions of
229 the mass of our Sun. Typically, a small galaxy contains a smaller black hole,
230 for example 50,000 times the mass of our Sun, while a large black hole, such as
231 the last entry in Table 1, has a mass billions of times the mass of our Sun.

?

232 **Objection 4.** *You are being totally inconsistent about mass! In Chapter 1*
233 *we heard about the mass m of a stone; there you said nothing about mass*
234 *in units of length. Now you define M with length units. Make up your mind!*

TABLE 3.1 Masses of some astronomical objects.

Object	Mass in kilograms	Geometric measure of mass	Equatorial r -coordinate
Earth	5.9742×10^{24} kilograms	4.44×10^{-3} meters or 0.444 centimeters	6.371×10^6 meters or 6371 kilometers
Sun	1.989×10^{30} kilograms	1.477×10^3 meters or 1.477 kilometers	6.960×10^8 meters or 696 000 kilometers
Black hole at center of our galaxy	8×10^{36} kilograms (4×10^6 Sun masses)	6×10^9 meters	
Black hole in galaxy NGC 4889	4.2×10^{40} kilograms (21×10^9 Sun masses)	3.1×10^{13} meters	



235
236
237
238
239
240
241
242
243

Excellent point. The difference between the mass M of a center of attraction and the mass m of a stone is important. First, a stone is a “free particle . . . whose mass warps spacetime too little to be measured” (inside the back cover). Second, most often we combine the stone’s mass m with another quantity in such a way that the result is a unitless ratio—for example E/m —by choosing the *same* unit in numerator and denominator. It does not matter which unit we use—joules or kilograms or electron-volts or the mass of the proton—as long as we use the *same* unit in numerator and denominator.

244
245
246
247
248

In contrast to the stone, the mass of a star or black hole *does* curve and warp spacetime. In this book the capital letter M *always* signals this fact. Here too we can arrange things so that M appears in a unitless ratio, such as $2M/r$, in which case M and r must have the same unit, which we choose to be meters.



249
250
251
252
253
254

Objection 5. *Okay, terrific, and this gives me a great idea: Why not simplify things even more by using unitless spacetime coordinates. Divide the Schwarzschild metric through by M^2 , then define dimensionless coordinates $\tau^* \equiv \tau/M$ and $t^* \equiv t/M$ and $r^* \equiv r/M$. Here the asterisk (*) reminds us that we are using dimensionless coordinates. Now the timelike Schwarzschild metric takes the simplest possible form:*

$$d\tau^{*2} = \left(1 - \frac{2}{r^*}\right) dt^{*2} - \frac{dr^{*2}}{\left(1 - \frac{2}{r^*}\right)} - r^{*2} d\phi^2 \quad (11)$$

(unitless coordinates)

255
256
257
258

*This notation has two big advantages: First, our equations are no longer cluttered with the symbol M , just as we have already eliminated from our equations the clutter of constants G and c . Second, metric (11) applies automatically to **all** black holes, of whatever mass M .*

3-10 Chapter 3 Curving

Box 4. “Our Little Jugged Apocalypse”

We tend to think of a black hole as a large object, especially the “monster” at the center of our galaxy (Table 1). But the word *large* invites the question, “Large compared to what?” The diameter of the black hole in our galaxy is about 10^{-6} light year. Our galaxy, a typical one, is some 10^5 light years in diameter. Any object a factor 10^{-11} the size of a galaxy must be considered a relative dot in the galactic scheme of things. Its relatively small size allows us to call the black hole

our “little juggled apocalypse,” a phrase the writer John Updike uses to describe the view into the portal of a front-loading clothes-washing machine. Conveniently, spacetime curvature increases from zero far from the isolated black hole to an unlimited value at its singularity. This makes the black hole a useful example to teach large swaths—but not all—of general relativity.



259
260
261
262
263
264
265
266
267
268

Originally we used your idea for a few chapters, but then returned these chapters to our current notation, which has several advantages: (1) Keeping the M allows us to check units in every equation. An equation can be wrong if the units are correct, but it is *always* wrong if the units are incorrect! (2) We can return to flat spacetime and special relativity simply by letting $M \rightarrow 0$; a second useful check. (3) We prefer to be continually reminded of the concrete *heft*—the observed massiveness—of astronomical objects: stars and black holes. For these reasons we choose to retain coordinates in units of length and the explicit symbol M in our equations.

Newton’s gravity
with mass in meters

269
270
271
272
273
274
275

How does Newton’s law of gravitation change when we express mass in meters? Think of a stone of mass m_{kg} near a center of attraction of mass M_{kg} . Rewrite Newton’s second law of motion ($F = ma$) for this case, using the gravitational force equation (7), with $m_{\text{kg}}g_{\text{conv}}$ on the left, where g_{conv} is the local acceleration of gravity. The stone’s mass m_{kg} cancels from both sides of the resulting equation. A minus sign signals that the acceleration is in the decreasing r direction.

$$g_{\text{conv}} = -\frac{GM_{\text{kg}}}{r^2} \quad (\text{Newton, conventional units}) \quad (12)$$

276
277

Now divide both sides of (12) by c^2 so as to obtain the conversion factor of equation (9). We can then write

$$g \equiv \frac{g_{\text{conv}}}{c^2} = -\frac{M}{r^2} \quad (\text{Newton, mass in meters}) \quad (13)$$

Newton’s g_{Earth}
with mass in meters

278
279
280
281
282

Remember that this is an equation of Newton’s mechanics, not an equation of general relativity. The quantities M and r both have the unit meter, so g has the unit meter^{-1} . Substitute into (13) the values of M_{Earth} and r_{Earth} from inside the front cover to obtain the value for the acceleration of gravity g_{Earth} at Earth’s surface in units of inverse meters:

$$g_{\text{Earth}} = -\frac{M_{\text{Earth}}}{r_{\text{Earth}}^2} = -1.09 \times 10^{-16} \text{ meter}^{-1} \quad (\text{Newton, mass in meters}) \quad (14)$$




FIGURE 3 US Pavillion “geodesic dome” designed by R. Buckminster Fuller for the 1967 International and Universal Exposition in Montreal. Place a clock at every intersection of rods on the outer surface of this sphere to create a small model of our imaginary nested spherical shells concentric to a black hole. Image courtesy of the Estate of R. Buckminster Fuller.

283 Does this numerical value seem small? It is the same acceleration we are used
 284 to, just expressed in different units. To jump from a high place on Earth is
 285 dangerous, whatever units you use to describe your motion!

286 Next we continue the explanation of Schwarzschild metrics (5) and (6)
 287 with a definition of the global radial coordinate r in these equations.

3.3 ■ THE GLOBAL SCHWARZSCHILD r -COORDINATE

289 *Measure the r -coordinate while avoiding the trap in the center*

Why Schwarzschild
 global coordinates?

290 Section 2.5 asked, “Does the black hole care what global coordinate system we
 291 use in deriving our global spacetime metric?” and answered, “Not at all!”
 292 General relativity allows us to use *any global coordinate system whatsoever*,
 293 subject only to some requirements of smoothness and uniqueness (Section 5.9).
 294 *Next question:* Since Schwarzschild had (almost) complete freedom to choose
 295 his global coordinates t, r , and ϕ , why did he choose the particular coordinates
 296 that appear in (5) and (6)? *Next answer:* Schwarzschild’s global coordinates
 297 take advantage of the spherical symmetry of a non-spinning black hole. When
 298 these coordinates are submitted to Einstein’s equations, they return metrics
 299 that are (relatively!) simple. In this and the following section we introduce and
 300 describe Schwarzschild global coordinates.

3-12 Chapter 3 Curving

Spherical shell
of rods and clocks

301 Start with Schwarzschild's r coordinate: Take the center of attraction to
302 be a black hole with the same mass as our Sun. In imagination, build around
303 it a spherical shell of rods fitted together in an open mesh (Figure 3). On this
304 shell mount a clock at every intersection of these rods. The rods and clocks of
305 such a collection of shells provides one system of coordinates to determine the
306 location of events that occur outside the event horizon.

We cannot measure
 r -coordinate directly.

307 How shall we define the size of the sphere formed by this latticework shell?
308 Shall we measure directly the radial separation between the sphere's surface to
309 its center? That won't do. Yes, in imagination we can stand on the shell. Yes,
310 we can lower a plumb bob on a "string." But for a black hole, any string, any
311 tape measure, any steel wire—whatever its strength—is relentlessly torn apart
312 by the unlimited pull the black hole exerts on any object that dips close
313 enough to its center. And even for Earth or Sun, the surface itself keeps us
314 from lowering our plumb bob directly to the center.

Derive r -coordinate
from measurement
of circumference.

315 Therefore try another method to define the size of the spherical shell.
316 Instead of lowering a tape measure from the shell, run a tape measure around
317 it in a great circle. The measured distance so obtained is the *circumference* of
318 the sphere. Divide this circumference by $2\pi = 6.283185\dots$ to obtain a distance
319 that would be the directly-measured r -coordinate of the sphere *if* the space
320 inside it were flat. But that space is *not flat*, as we shall see. Yet this procedure
321 yields the most useful known measure of the size of the spherical shell.

Definition:
 r -coordinate

322 The "radius" of a spherical object obtained by this method of measuring
323 has acquired the name **r -coordinate**, because it is no genuine Euclidean
324 radius. We call it also the **reduced circumference**, to remind us that it is
325 derived ("reduced") from the circumference:

$$\begin{aligned} r\text{-coordinate} &\equiv \text{reduced circumference} && (15) \\ &\equiv \frac{\text{measured circumference}}{2\pi} \end{aligned}$$

326 We sometimes use the expression Schwarzschild- r , which labels the global
327 coordinate system of which r is a member. From now on we try not to use the
328 word "radius" for the r -coordinate, because it can confuse results for flat
329 spacetime with results for curved spacetime.

330 During construction of each shell the contractor stamps the value of its
331 r -coordinate on it for all to see.

?

332 **Objection 6.** *Aha, gottcha! To define the r -coordinate in (15), you*
333 *measure the length of the entire circumference of a spherical shell. Near a*
334 *massive black hole, this circumference could be hundreds of kilometers*
335 *long. Yet from the beginning you say, "Report every measurement using a*
336 *local inertial frame." Near a black hole a local inertial frame is tiny*
337 *compared with the length of this circumference. You do not follow your own*
338 *rules for measurement.*

Section 3.3 The Global Schwarzschild r -coordinate **3-13**




FIGURE 4 The scale of some objects described by physics. Objects close to the diagonal line are those for which correct predictions require general relativity. See Box 5. Figure adapted from the textbook *Gravity* by James Hartle.

339
340

Guilty as charged! We failed to spell out the process: Use a whole string of overlapping local inertial frames parked around the circumference of the

3-14 Chapter 3 Curving

Box 5. When is General Relativity Necessary?

When is general relativity required to describe and predict accurately the behavior of structures and phenomena in our Universe? See Figure 4.

ORDINARY STAR. An ordinary star like our Sun does not require general relativity to account for its development, structure, or physical properties. Like all massive centers of attraction, however, it does deflect and focus passing light in ways accounted for by general relativity (Chapter 13).

WHITE DWARF. A white dwarf is the burned out cinder of an ordinary star, with a mass approximately equal to that of our Sun and r -coordinate of its surface comparable to that of Earth. General relativity is not required to account for the structure of the white dwarf but is needed to predict stability, especially near the so-called **Chandrasekhar limit** of mass—about 2.4 times the mass of our Sun—above which the white dwarf is doomed to collapse.

NEUTRON STAR. A neutron star can result from the collapse of a white dwarf star. Its mass is approximately that of our

Sun with an r -coordinate of its surface about 10 kilometers, the size of a city. General relativity significantly affects the structure and oscillations of the neutron star. Emission of gravitational waves (Chapter 16) may damp out non-radial vibrations.

BLACK HOLE. “The physics of black holes calls on Einstein’s description of gravity from beginning to end.” (Misner, Thorne, and Wheeler)

GRAVITY WAVES. We have observed gravitational radiation predicted by general relativity.

THE UNIVERSE. Models of the Universe as a single structure employ general relativity (Chapters 14 and 15). It seems increasingly likely that general relativity correctly accounts for non-quantum features of the Universe, but it remains possible that general relativity fails over these immense spans of spacetime and must be replaced by a more general theory.

341 spherical shell, then define the circumference to be the summed measured
 342 distances across each of these local inertial frames. In practice this
 343 procedure is awkward, but in principle it avoids your otherwise valid
 344 objection.

Directly-measured
 separation between
 nested shells is **greater**
 than the difference in
 their r -values.

345 Think of building two concentric shells, a lower shell of reduced
 346 circumference r_L and a higher shell of reduced circumference r_H , such that the
 347 difference in reduced circumference $r_H - r_L$ equals 100 meters. Stand on the
 348 higher shell and lower a plumb bob, and for the first time measure directly the
 349 radial separation perpendicularly from the higher shell to the lower one. Will
 350 we measure a 100-meter radial separation between our two shells? We would if
 351 space were flat. But outside a massive body space is *not* flat. The relation
 352 between global differential dr and measured radial differential $d\sigma$ comes from
 353 the spacelike version of the Schwarzschild metric (6) with $dt = d\phi = 0$.

$$d\sigma = \frac{dr}{\left(1 - \frac{2M}{r}\right)^{1/2}} \quad (\text{radial shell separation, } dt = d\phi = 0) \quad (16)$$

354 We note immediately that for the radial shell separation $d\sigma$ to be a real
 355 quantity, we must have $r > 2M$; otherwise the square root in the denominator
 356 has an imaginary value. This is an indication that shells can be built only
 357 outside the event horizon (Section 6.7).

358 Outside the event horizon, the magnitude of the denominator on the right
 359 side of (16) is always less than one. Hence Schwarzschild geometry tells us that

Section 3.3 The Global Schwarzschild r -coordinate 3-15

360 every radial differential increment $d\sigma$ is *greater* than the corresponding
 361 differential increment dr of the reduced circumference. Therefore the summed
 362 (integrated) measureable distance between our two shells is greater than 100
 363 meters, even though their circumferences differ by exactly $2\pi \times 100$ meters. This
 364 discrepancy between measured separation and difference in global r -coordinate
 365 provides striking evidence for the curvature of space. See Sample Problem 1.

Small effect
near our Sun

366 Built around our Sun, the r -coordinate of the inner shell cannot be less
 367 than that of our Sun's surface, 695 980 kilometers. Around this inner shell we
 368 erect a second one—again in imagination—of r -coordinate 1 kilometer greater:
 369 695 981 kilometers. The directly-measured distance between the two would be
 370 not 1 kilometer, but 2 millimeters *more* than 1 kilometer.

Get closer
to the center.

371 How can we get closer to the center of a stellar object with mass equal to
 372 that of our Sun—but still remain external to the surface of that object? A
 373 white dwarf and a neutron star each has roughly the same mass as our Sun,
 374 but each is much smaller than our Sun. So we can—in principle—conduct a
 375 more sensitive test of the nonflatness of space much closer to the centers of
 376 these objects while staying external to them (Box 5). The effects of the
 377 curvature of space are much greater near the surface of a white dwarf than near
 378 the surface of our Sun—and greater still near the surface of a neutron star.

?

379 **Objection 7.** *Why not define the r -coordinate differently—call it r_{new} —in*
 380 *terms of the directly-measured distance between two adjacent shells. For*
 381 *example, we could give the innermost shell at the event horizon the radial*
 382 *coordinate $r_{\text{new}} = 2M$, and the next shell $r_{\text{new}} = 2M + \Delta\sigma$, where $\Delta\sigma$*
 383 *is the directly-measured separation between that shell and the innermost*
 384 *shell. And so on. That would eliminate the awkwardness of your quoted*
 385 *results.*

!

386 You can choose (almost) any global coordinate system you want, but the
 387 one you suggest is inconvenient. First, you cannot escape the deviation
 388 from Euclidean geometry imposed by curvature; your definition leads to a
 389 calculated circumference $2\pi r_{\text{new}}$ that is different from the
 390 directly-measured one. Second, outside the event horizon your definition is
 391 awkward to carry out, since it requires collaboration between observers on
 392 different shells. Third, how is your definition applied inside the event
 393 horizon, where no shells exist? (Box 7 in Section 7.8 shows how to
 394 measure the Schwarzschild reduced circumference r inside the event
 395 horizon.) Finally, your definition of r_{new} , when submitted with t and ϕ to
 396 Einstein's equations, results in a different metric—a more complicated
 397 one—which would be more inconvenient to use than the Schwarzschild
 398 global metric.

Huge effect
near black hole

399 Turn attention now to a black hole of mass M . Close to it the departure
 400 from flatness is much larger than it is anywhere around a white dwarf or a
 401 neutron star. Construct an inner shell having an r -coordinate, a reduced
 402 circumference, of $3M$. Let an outer shell have an r -coordinate of $4M$. In
 403 contrast to these two r -coordinates, defined by measurements around the two
 404 shells, the directly-measured radial distance between the two shells is $1.542M$,

3-16 Chapter 3 Curving

Sample Problem 1. “Space Stretching” Near a Black Hole

Here we verify the statement near the end of Section 3.3 that for a black hole of mass M , the directly-measured radial distance calculates as $1.542M$ between the lower shell at r -coordinate $r_L = 3M$ and the higher shell at r -coordinate $r_H = 4M$. In Euclidean geometry this measured distance would be $1.000M$, but not in curved space!

SOLUTION Equation (16) gives the radial differential $d\sigma$ between shells separated by a differential dr of the global radial coordinate r . The term $2M/r$ changes significantly over the range from $r = 3M$ to $r = 4M$, so our “summation” must be an integral. Integrating (16) from lower r -coordinate $r_L = 3M$ to higher r -coordinate $r_H = 4M$ yields:

$$\begin{aligned} \sigma &= \int_{r_L}^{r_H} \frac{dr}{\left(1 - \frac{2M}{r}\right)^{1/2}} \\ &= \int_{r_L}^{r_H} \frac{r^{1/2} dr}{(r - 2M)^{1/2}} \end{aligned} \quad (17)$$

This integral is not in a common table of integrals, so make the substitution $r = z^2$, from which $dr = 2zdz$. The resulting integral has the solution:

$$\begin{aligned} \sigma &= \int_{z_L}^{z_H} \frac{2z^2 dz}{(z^2 - 2M)^{1/2}} \\ &= \left[z(z^2 - 2M)^{1/2} + 2M \ln \left| z + (z^2 - 2M)^{1/2} \right| \right]_{z_L}^{z_H} \end{aligned} \quad (18)$$

Here the symbol \ln (spelled “ell” “en”) represents the natural logarithm (to the base e) and vertical-line brackets indicate absolute value. Substitute the values

$$z_L = (3M)^{1/2} \quad \text{and} \quad z_H = (4M)^{1/2}$$

and recall that for logarithms, $\ln(B) - \ln(A) = \ln(B/A)$. The result is

$$\sigma = 1.542M \quad (\text{radial, exact}) \quad (19)$$

Here the symbol σ predicts the *exact* radial separation between these shells measured by the shell observer who uses a short ruler, say one-centimeter long, laid end to end many times to find a total measured distance. This exact result is radically different from $1.000M$ predicted by Euclid.

405 compared to the Euclidean-geometry figure of $1.000M$ (Sample Problem 1). At
 406 this close location, the curvature of space results in measurements quite
 407 different from anything that textbook Euclidean geometry would lead us to
 408 expect. We call this effect the **stretching of space**.



409 **Objection 8.** *WHY is the directly-measured distance between spherical*
 410 *shells greater than the difference in r coordinates between these shells? Is*
 411 *this discrepancy caused by gravitational stretching or compression of the*
 412 *measuring rods?*



413 No, the quoted result assumes rigid measuring equipment. In practice, of
 414 course, a measuring rod held by the upper end will be subject to
 415 gravitational stretching (or compression if held by the lower end). Make the
 416 rod short enough; then gravitational stretching is unimportant. Now count
 417 the number of times the rod has to be moved end to end to cross from one
 418 shell to the other.



419 **Objection 9.** *Are you refusing to answer my question? What CAUSES the*
 420 *discrepancy, the fact that the directly-measured distance between*
 421 *spherical shells is greater than the difference in r coordinates between*
 422 *these shells? WHY this discrepancy?*

Sample Problem 2. Our Sun Causes Small Curvature

The Schwarzschild metric function $(1 - 2M/r)$ gauges the difference between flat and curved spacetime. How far from the center of our Sun must we be before the resulting curvature becomes extremely small or negligible?

A. As a first example, find the r -coordinate from a point mass with the mass of our Sun ($M \approx 1.5 \times 10^3$ meters) such that the metric function differs from the value one by one part in a million. Compare this r -coordinate to the actual r -coordinate of the surface of our Sun ($r_{\text{Sun}} \approx 7 \times 10^8$ meters).

B. As a second example, find the radial r -coordinate from our Sun such that the metric function differs from the value one by one part in 100 million. Compare the value of this r -coordinate with the average r -coordinate of Earth's orbit ($r \approx 1.5 \times 10^{11}$ meters).

SOLUTIONS

A. We want $(1 - 2M/r) \approx 1 - 10^{-6}$, which yields

$$r \approx \frac{2M}{10^{-6}} = 2 \times 1.5 \times 10^3 \times 10^6 \text{ meters} \quad (20)$$

$$= 3 \times 10^9 \text{ meters}$$

This r -coordinate is approximately four times the r -coordinate of our Sun's surface.

B. In this case we want $(1 - 2M/r) \approx 1 - 10^{-8}$, so

$$r \approx \frac{2M}{10^{-8}} = 2 \times 1.5 \times 10^3 \times 10^8 \text{ meters} \quad (21)$$

$$= 3 \times 10^{11} \text{ meters}$$

which is approximately twice the r -coordinate of Earth's orbit.

423 
 424
 425
 426
 427
 428
 429

A deep question! Fundamentally, this discrepancy shatters the notion of Euclidean space. We are faced with a weird measured result, which we can summarize with the statement, "Mass stretches space." Your question "Why?" is not a scientific question, and science cannot answer it. We know only observed results and their derivation from general relativity. Does the following satisfy you? *Space stretching causes the discrepancy!* Section 3.8 exhibits one way to visualize this stretching.

3.4 THE GLOBAL SCHWARZSCHILD t -COORDINATE

431 *Freeze global space coordinates; examine the warped t -coordinate.*

To describe orbits,
we need curvature
of spacetime.

432 It is not enough to know the results of curvature on the r -coordinate alone. To
 433 appreciate how the grip of spacetime tells planets how to move requires us to
 434 understand how curvature affects the global t -coordinate as well. The
 435 coordinate differential dt appears on the right side of the Schwarzschild metric.
 436 Basically, Schwarzschild's definition of the t -coordinate was arbitrary, like the
 437 definition of every global coordinate.

Relation between
 $d\tau$ and dt

438 How does Schwarzschild coordinate differential dt relate to the differential
 439 wristwatch time $d\tau$ between two successive events that occur at fixed r - and
 440 ϕ -coordinates? The coordinate differentials dr and $d\phi$ are both equal to zero
 441 for that pair of events. Then the interval between ticks is the wristwatch time
 442 derived from metric (5), that is:

$$d\tau = \left(1 - \frac{2M}{r}\right)^{1/2} dt \quad (\text{stationary clock: } dr = d\phi = 0) \quad (22)$$

443 Equation (22) shows that far from a black hole ($r \rightarrow \infty$), Schwarzschild- t
 444 coincides with the time of a shell clock located there. This is an important,

3-18 Chapter 3 Curving

445 but accidental, convenience of Schwarzschild's choice of global t -coordinate. It
 446 is not true for the metrics of many other global coordinate systems for the
 447 non-spinning black hole.

Slogan:
 "A clock at
 smaller r
 runs slower."

448 Now look at the prediction of equation (22) closer to a black hole—but
 449 still outside the event horizon. There the Schwarzschild coordinate differential
 450 dt will be *larger* than the differential wristwatch time $d\tau$ measured by a clock
 451 at rest on the shell at that r -coordinate. Smaller wristwatch time $d\tau$ between
 452 two standard events leads to the useful but somewhat imprecise slogan, *A*
 453 *clock closer to a center of attraction runs slower* (see Section 4.3).

Schwarzschild:
 complete description

454 We have now carefully defined each of the Schwarzschild global coordinates
 455 and displayed the resulting global metric handed to us by Einstein's equations,
 456 including the range of global coordinates given in equations (5) and (6). This
 457 combination—plus its connectedness (topology)—provides a *complete*
 458 description of spacetime near the isolated non-spinning black hole. These tools
 459 alone are sufficient to determine every (classical, that is non-quantum)
 460 observable property of spacetime in this region.

?

461 **Objection 10.** *Hold it! You gave us separate Sections 3.3 and 3.4 on two*
 462 *global coordinates, Schwarzschild- r and Schwarzschild- t , respectively.*
 463 *Why no section on the third global coordinate, Schwarzschild- ϕ ?*

!

464 Good question. In answer, compare metric (4) for flat spacetime in Box 1
 465 with the Schwarzschild metric (5) for curved spacetime. The last term is
 466 the same in both equations: $-r^2 d\phi^2$. Typical in relativity, the t -coordinate
 467 gives us the most trouble and the r -coordinate less trouble. In the
 468 non-spinning black hole metrics used in this book, the angle ϕ gives no
 469 trouble at all, due to the angular symmetry. For the spinning black hole
 470 (Chapters 17 through 21), however, even this angle becomes a
 471 troublemaker!

3.5. ■ CONSTRUCTING THE GLOBAL SCHWARZSCHILD MAP OF EVENTS

473 *Read a road map, but don't drive on it!*

"Think globally;
 measure locally."

474 In this book we choose to make every measurement and observation in a local
 475 inertial frame. But that does not suffice to describe the relation between
 476 events far from one another in the vicinity of the black hole. Suppose we know
 477 the stone's energy and momentum measured in one local inertial frame
 478 through which it passes. How can we predict the stone's energy and
 479 momentum in a second local inertial frame far from the first?

480 This prediction requires (a) knowledge of the stone's initial location in
 481 global coordinates, (b) analysis of the global worldline of the stone between
 482 widely-separated local frames, and (c) conversion of a piece of the global
 483 trajectory to local inertial coordinates in the remote inertial frame. This
 484 section begins that process, which we summarize with the slogan "*Think*
 485 *globally, measure locally.*"

Section 3.5 Constructing the Global Schwarzschild Map of Events 3-19




FIGURE 5 *Schwarzschild map of the trajectory of a free stone that falls into a black hole. As it falls, it emits (numbered) flashes equally separated in time on its wristwatch. However, these flash emissions are not equally spaced along the Schwarzschild map trajectory. Each numbered event also has its Schwarzschild- t . NO ONE observes directly the entire trajectory shown on this map. Question: Why are numbered emission events closer together near both ends of the trajectory than in the middle of the trajectory? The answer for events 1 through 3 should be simple. The answer for events 5 through 8 appears in Section 6.5.*

486 Global Schwarzschild coordinates locate events around a black hole similar
 487 to the way in which latitude and longitude locate places on Earth’s surface
 488 (Section 2.3). A global map of Earth is nothing but a rule that assigns unique
 489 coordinates to each *point* on its surface.

The **spacetime map**
 assigns coordinates
 to every event.

490 By analogy, we speak of a **spacetime map**, which is nothing but a rule
 491 that assigns unique coordinates to each *event* in the region described by that
 492 map. This section describes the construction and uses of the Schwarzschild
 493 spacetime map, a task that we personalize as the work of an archivist.

**Schwarzschild
 mapmaker**

494 Think of Schwarzschild coordinates as an accounting system, a
 495 bookkeeping device, a spreadsheet, a tabulating mechanism, an international
 496 language, a space-and-time database created by an archivist who records every
 497 event and all motions in the entire spacetime region exterior to the surface of
 498 the Earth or Moon or Sun—or anywhere around a black hole except exactly at
 499 its center. We personify the supervisor of this record as the **Schwarzschild
 500 mapmaker**. The Schwarzschild mapmaker receives reports of actual
 501 measurements made by local shell and other inertial observers, then converts
 502 and combines them into a comprehensive description of events (in
 503 Schwarzschild coordinates) that spans spacetime around a black hole. The
 504 mapmaker makes no measurements himself and does not analyze
 505 measurements. He is a data-handler, pure and simple.

Mapmaker:
 the central
 coordinator.

506 The Schwarzschild mapmaker (or his equivalent) is absolutely necessary
 507 for a complete description of the motion of stones and light signals around a
 508 black hole. He has the central coordinating role in describing globally all the
 509 events that take place outside the event horizon of the black hole. He collates

3-20 Chapter 3 Curving

Box 6. The Metric as Spacetime Micrometer



FIGURE 6 The micrometer caliper measures directly a tiny distance or thickness, bypassing x and y coordinates. The watch measures directly the invariant wristwatch time between two events, bypassing separate global coordinate increments. (Photo by Per Torphammar.)

What *is* the metric? What is it *good for*? Think of a **micrometer caliper** (Figure 6), a device used by metalworkers and other practical workers to measure a small distance. The micrometer caliper translates turns of a calibrated screw on the cylinder into the directly-measured distance across the gap between the flat ends of the little cylinders in the upper right corner of the figure. The worker *owns* the micrometer; the worker *chooses* which distance to measure with the micrometer caliper.

The metric is our “four-dimensional micrometer” that translates global coordinate separations between an adjacent pair of events into the measurable wristwatch time lapse or ruler distance between those events. You *own* the metric. You *choose* the events whose separation you wish to measure with the metric.

1. **One possible choice:** Two sequential ticks of a clock bolted to a spherical shell. Then $dr = d\phi = 0$ and the

wristwatch time $d\tau$ is the time lapse read directly on the shell clock.

2. **A second possible choice:** Events with the same global t -coordinate that occur at the two ends of a stick held at rest radially between two adjacent shells, so that $dt = d\phi = 0$. Then the ruler distance $d\sigma$ is the directly-measured length of the stick—equation (16).
3. **A third possible choice:** Two sequential ticks on the wristwatch of a stone in free fall along a radial trajectory. Then $d\phi = 0$ and $d\tau$ is read directly on the wristwatch.

And so on. There are an infinite number of event-pairs near one another that you can choose for measurement using your four-dimensional micrometer—the metric.

Assembling many micrometer caliper measurements can in principle describe the geometry of space. Assembling many wristwatch and ruler measurements can in principle describe the geometry of spacetime: “The metric completely specifies local spacetime and gravitational effects within the global region in which it applies.” (Inside back cover.)

What advice will the “old spacetime machinist” give to her younger colleague about the practical use of the metric? She might share the following pointers:

1. Focus on *events* and the separation between each pair of events, not fuzzy concepts like “time” or “location.”
2. Do not confuse results from one pair of events with results from another pair of events.
3. Whenever possible, choose two adjacent events for which the increment of one or more map coordinates is zero.
4. Whenever possible, identify the wristwatch time or ruler distance with some observer’s direct measurement.
5. When a light flash moves directly from one event to another event, the wristwatch time *and* the ruler distance between those events are both zero: $d\tau = d\sigma = 0$.

510 data from many local observers and combines them in various ways, for
511 example drawing a global map such as the one plotted in Figure 5.

512 The Schwarzschild mapmaker can be located anywhere. How does he learn
513 of events in his dominion? Like a taxi dispatcher, he uses radio to keep track of
514 moving stones, light flashes, and in addition locates explosions and other
515 events of interest, perhaps as follows:

516 Stamped on each spherical shell is its map r -coordinate; we mark different
517 locations around the shell with different values of ϕ . At each location place a
518 recording clock that reads the Schwarzschild- t (Box 6). Each clock radios to

Box 7. Where does the event horizon come from?

The event horizon—that one-way spacetime surface that lets light and stones pass inward but forbids them to cross outward—is a surprise. Who could have predicted it? Answer: Nobody did.

Newton readily predicts the gravitational consequences of a point mass, telling us immediately the initial acceleration of a stone released from rest at any r -coordinate. Twice the attracting mass, twice the stone's acceleration at that r -value; a million times the attracting mass, a million times the stone's acceleration. Newton's theory of gravity is *linear* in mass.

Not so for Einstein's general relativity, which is relentlessly *nonlinear*. In general relativity not only mass but also energy and pressure curve spacetime. A star of twice the mass typically has increased internal pressure, resulting in more than twice the gravitational effects at the same r -coordinate outside its surface. For an ordinary star the added effect of

pressure is negligible; for a neutron star the added effect of pressure is important; for a black hole the added effect of pressure is catastrophic.

When a neutron star, for example, steals mass from a normal companion star, the pressure near its center increases, along with the added matter. The net result is greater than that due to the added matter alone. At a certain point, this process "runs away," resulting in collapse into a black hole.

Linear effects mean proportional response in phenomena. Nonlinear effects lead to entirely new phenomena. For the non-spinning black hole, a major outcome of nonlinearity is the event horizon. Near to the *spinning* black hole (Chapters 17 through 21), the nonlinearity of Einstein's theory leads to an even more complex geometry of spacetime and consequent radical, unexpected physical effects.

Mapmaker: top level, bureaucrat

519 the mapmaker the nature of an event that occurs next to it, along with its
520 global coordinates (t, r, ϕ) . After inevitable transmission delays due to the
521 finite speed of light, the mapmaker at the control center assembles a global
522 Schwarzschild map that gives coordinates and description of every
523 measurement and observation. Our mapmaker acts as a top-level bureaucrat.

Using the Schwarzschild map

524 No one lives on a road map, but we use it to describe the territory and to
525 plan our trip. Similarly, coordinates r, ϕ , and t are simply labels on a spacetime
526 map. These coordinates uniquely locate events in the entire spacetime region
527 outside the surface of any spherically symmetric gravitating body or anywhere
528 around a black hole except on its singularity. The Schwarzschild map guides
529 our navigation near a black hole, in the same way that an arbitrary set of
530 global coordinates—made into maps—guides our travels on Earth's surface.

Map coordinate difference \neq measured length or time lapse.

531 But never forget: In most cases Schwarzschild map coordinate separations
532 are *not* what any local inertial observer measures directly.

533 *Advice: It is best never to confuse a global map coordinate separation with*
534 *the local inertial frame measurement of a distance or time lapse.* More details
535 in Chapter 5.



536 **Objection 11.** *Stop giving me second-hand ideas! I want **reality**. Your*
537 *concept of a Schwarzschild map is nothing but an analogy to the inevitable*
538 *distortions in geography when Earth's spherical surface is squashed onto*
539 *a flat map. Where is the true representation of curved spacetime,*
540 *corresponding to the true spherical map of Earth's surface?*



541 Early in the history of sea travel, mapmakers thought the world was flat. An
542 ancient sea captain acquainted with Euclid's plane geometry (and also the

3-22 Chapter 3 Curving

543 much later calculus differential notation of Leibniz!) would puzzle over the
 544 metric for differential distance ds on Earth's surface, equation (3) in
 545 Section 2.3:

$$ds^2 = R^2 \cos^2 \lambda d\phi^2 + R^2 d\lambda^2 \quad (\text{space metric: Earth's surface}) \quad (23)$$

546 The ancient sea captain asks, "What is R ?" (r -coordinate of the Earth's
 547 surface). "What are λ and ϕ ?" (angles of latitude and longitude). "Why
 548 does differential distance ds depend on latitude λ ?" (convergence at the
 549 poles of lines of constant longitude). "Where is the edge?" (There is no
 550 edge.) Who is responsible for the captain's perplexity about a curved
 551 surface? Not Nature; not Mother Earth. Neither is Nature responsible for
 552 our perplexity about curved spacetime. Everything will be crystal clear as
 553 soon as we can visualize four-dimensional curved spacetime. But we do
 554 not know anyone who can do this; we certainly cannot! So we
 555 compromise, we do our best to live with our limitations and to develop
 556 intuition from the analogy to curved surfaces in space, such as the partial
 557 visualization of Schwarzschild geometry in the following sections.

558 **Black holes just didn't "smell right"**

559 *During the 1920s and into the 1930s, the world's most renowned experts*
 560 *on general relativity were Albert Einstein and the British astrophysicist*
 561 *Arthur Eddington. Others understood relativity, but Einstein and*
 562 *Eddington set the intellectual tone of the subject. And, while a few others*
 563 *were willing to take black holes seriously, Einstein and Eddington were*
 564 *not. Black holes just didn't "smell right"; they were outrageously bizarre;*
 565 *they violated Einstein's and Eddington's intuitions about how our*
 566 *Universe ought to behave . . . We are so accustomed to the idea of black*
 567 *holes today that it is hard not to ask, "How could Einstein be so dumb?*
 568 *How could he leave out the very thing, implosion, that makes black*
 569 *holes?" Such a reaction displays our ignorance of the mindset of nearly*
 570 *everybody in the 1920s and 1930s . . . Nobody realized that a sufficiently*
 571 *compact object must implode, and that the implosion will produce a black*
 572 *hole.*

573 —Kip Thorne

3.6. ■ THE SPACETIME SLICE

575 *Do the best we can to visualize curved spacetime*

576 This section introduces a method of visualizing curved spacetime—called the
 577 **spacetime slice**—that we use repeatedly throughout the book. Every such
 578 visualization of curved spacetime is partial and incomplete—it does not tell
 579 all—but can carry us some of the way toward intuitive understanding of
 580 spacetime curvature.

581 **DEFINITION 2. Spacetime slice**

582 A **spacetime slice**—which we usually just call a **slice**—is a
 583 two-dimensional spacetime surface on which we plot two global
 584 coordinates of all events that lie on that surface and that have equal




FIGURE 7 *Preview:* When we apply the global metric to a slice, then on every region of the slice we can either draw a light cone diagram or construct an embedding diagram.

Definition:
spacetime slice

585
586
587
588
589
590
591

values for all other global coordinates. We indicate a slice with square brackets; the three alternative slices for our Schwarzschild global coordinates are $[r, \phi]$, $[r, t]$, and $[\phi, t]$. Our definition of *slice* includes its range of coordinates and its connectedness (topology). The slice—even when populated with events—does not use the metric, so a *spacetime slice carries no information whatsoever about spacetime curvature*. This feature makes the slice useful in both special and general relativity.

On every region of every slice: light cone diagram or embedding diagram

592
593
594
595
596

The following remarkable property of the spacetime slice will illuminate the remainder of this book: *When we apply the global metric to a spacetime slice, then on every region of every slice we can either draw worldlines or set up an embedding diagram*. Figure 7 previews the content of the following sections.

597
598
599
600
601

What does “every region” of the slice mean in the caption to Figure 7? For the non-spinning black hole the regions are outside and inside the event horizon. Section 3.7 shows that light cones can be drawn on both regions for the $[r, t]$ slice. Section 3.9 shows that outside the event horizon the $[r, \phi]$ slice is an embedding diagram.

3-24 Chapter 3 Curving




FIGURE 8 Schwarzschild light cone diagram on an $[r, t]$ slice, constructed from segments of light worldlines from equation (26), showing future (F) and past (P) of each event (filled dots). At each r -coordinate the light cone can be moved up or down vertically without change of shape, as shown.

3.7.2 ■ LIGHT CONE DIAGRAM ON AN $[r, t]$ SLICE

603 *The global t -coordinate can run backward along a worldline!*

On an $[r, t]$ slice. . .

604 We can learn a lot about predictions of the Schwarzschild metric by plotting
 605 light cones. To derive the worldline of a light flash in r, t coordinates, set
 606 $d\tau = 0$ and $d\phi = 0$ in (5). The result is:

$$0 = \left(1 - \frac{2M}{r}\right) dt^2 - \left(1 - \frac{2M}{r}\right)^{-1} dr^2 \quad (\text{light, and } d\phi = 0) \quad (24)$$

607 Which leads to the equation:

$$\frac{dt}{dr} = \pm \frac{r}{r - 2M} \quad (\text{light, radial motion}) \quad (25)$$

Light cones

608 Integrate this to find the equation for the worldline of a light flash:

Box 8. A White Hole?




FIGURE 9 Schematic of a light cone inside the event horizon in Schwarzschild global coordinates.

Inside the event horizon, do a stone and light flash really move only toward smaller r ? And does Figure 8 correctly represent this? Why do the light cones not open upward in this figure, as they do in flat spacetime and also outside the event horizon?

To answer these questions, assume that the worldline of the stone passes through event E, the intersection of the light cone worldlines in Figure 9. Then determine what worldlines through E are possible between A and C (solid line) or between D and B (dashed line). The metric tells us how the stone's wristwatch advances along its constant- ϕ worldline. From (24), it reads

$$d\tau^2 = \left(1 - \frac{2M}{r}\right) dt^2 - \left(1 - \frac{2M}{r}\right)^{-1} dr^2 \quad (27)$$

Wristwatch time in (27) is real, therefore physical, only if the right side is positive. You can show that along a worldline connecting events D and B (dashed line), the wristwatch time is imaginary. In contrast, you can show that along a worldline that connects events A and C (solid line), wristwatch time is real. *First conclusion:* Worldlines of stones that pass through

event E can pass only from either the A region to the C region or from the C region to the A region. No stone worldline through event E can connect events B and D.

Next question: In which direction does the stone move between events A and C inside the event horizon? Arrows on the light cone imply that the motion is from A to C, namely to smaller r . But all differentials in (27) are squared: The metric allows motion in either direction.

We now show that motion to larger r cannot occur inside the event horizon. This means that the solution of the metric that allows motion to larger r inside the event horizon is an extraneous solution and does not correspond to the workings of Nature.

Suppose that the stone moves to larger r , from event C to event A, in which case the light cone arrows in Figure 9 would point to the right. That means that at an earlier wristwatch time the stone was at C. Now draw a light cone that crosses at event C. Then there is a still earlier event to the left of C through which the stone passed. Repeat this process until we reach $r = 0$, from which this stone must have emerged. The result is what we call a **white hole**. A white hole spews stones and light outward from its singularity, the opposite of a black hole.

Do white holes exist in Nature? We have not detected any. And if they should temporarily form, how could they possibly survive, since their central feature is to empty themselves into surrounding spacetime? The method we use here is called *reductio ad absurdum*, reduction to an absurd result.

Final conclusion: Arrows on the light cones inside the event horizon in Figure 9 point in the physically correct direction, which funnels stones and light toward the singularity. The corresponding light cones in Figure 8 do the same.

$$t - t_1 = \pm \left(r - r_1 + 2M \ln \left| \frac{r/M - 2}{r_1/M - 2} \right| \right) \quad (\text{light, radial motion}) \quad (26)$$

609 where (r_1, t_1) are initial coordinates of the light flash. Figure 8 plots the
610 resulting light cone diagram for many different values of (r_1, t_1) .

611 Figure 8 tells us a lot about physical predictions of the Schwarzschild
612 metric. The light cone of an event tells us the past (P) and future (F) of that
613 event. Note, first, that at the event horizon light does not change r -coordinate
614 on this slice. Second, inside the event horizon everything moves to smaller r .
615 The light cone corrals possible worldlines of a stone that passes through that

Trouble at the event horizon

3-26 Chapter 3 Curving

616 event—such as worldlines for Stone A and Stone B in the plot. Note, third,
617 that the t -coordinate runs backward along worldlines B and D.

?

618 **Objection 12.** *How can light be stuck at the event horizon, moving neither*
619 *inward or outward?*

!

620 Figure 8 tells us that near the event horizon the t -coordinate changes very
621 rapidly along a light ray, while the r coordinate changes very little. This is a
622 problem with the Schwarzschild t coordinate that obscures observed
623 results. We can say that the Schwarzschild t -coordinate is *diseased*, does
624 not correctly predict observations. Chapters 6 and 7 analyze and
625 overcome this global coordinate difficulty and show that light can fall to
626 smaller r , but not move to larger r inside the event horizon.

?

627 **Objection 13.** *Oops! How can time run backward along a worldline, such*
628 *as that of Stone B in Figure 8? Its arrow tends downward with respect to*
629 *the t/M axis.*

!

630 Careful! Never use the word “time” by itself (Section 2.7). Only the global
631 t -coordinate runs backward along worldlines B and D in Figure 9. Global
632 coordinates are (almost) totally arbitrary; we choose them freely, so we
633 cannot trust them to tell us what we will observe. Only the left side of the
634 metric does that, for example giving us wristwatch time between two
635 events. The wristwatch time is positive as the stone progresses along
636 worldline B in Figure 8; and along the worldline of *every* light flash the
637 wristwatch time is zero. Box 8 shows that the motion of both light and
638 stones must be to smaller r inside the event horizon.

?

639 **Objection 14.** *Aha! I've caught you in a serious contradiction. Inside the*
640 *horizon the worldline of the stone in Figure 8 is flatter than that of light.*
641 *That is, the stone traverses a greater span of r coordinate per unit time*
642 *than light does. The stone moves faster than light! Let's see you wiggle out*
643 *of that one!*

!

644 Again you use the word “time” incorrectly and compound the error by
645 changing r rather than moving a distance. Global coordinates are
646 arbitrary—our choice!—and global coordinate separations are not
647 measured quantities. This arbitrariness combines with spacetime
648 curvature to create the distortions plotted in Figure 8. Different global
649 coordinates give different distortions—see the same plot with different
650 global coordinates in Figure 5, Section 7.6. For *every* global coordinate
651 system dr/dt inside the event horizon does not *measure* the velocity of
652 anything. We favor measurement and observation on a local flat patch,
653 where special relativity rules. Chapter 5 has a lot more on this subject.

Section 3.8 Inside the Event Horizon: A Light Cone Diagram on an $[r, \phi]$ Slice **3-27**

3.8. ■ INSIDE THE EVENT HORIZON: A LIGHT CONE DIAGRAM ON AN $[r, \phi]$ SLICE

655 *Inside the event horizon, Schwarzschild- r is timelike!*

On an $[r, \phi]$ slice. . .

656 To continue our attempt to visualize curved spacetime around a black hole, we
657 plot light cones on an $[r, \phi]$ slice. Light plots on this slice require that $d\tau = 0$
658 and $dt = 0$. With these conditions, (5) becomes

$$0 = - \left(1 - \frac{2M}{r} \right)^{-1} dr^2 - r^2 d\phi^2 \quad (\text{light, and } dt = 0) \quad (28)$$

659 So the trajectory of light on the $[r, \phi]$ slice satisfies the equation:

$$\frac{d\phi}{dr} = \pm \frac{1}{r^{1/2}(2M - r)^{1/2}} \quad (\text{light, } dt = 0) \quad (29)$$

Light cones
inside the
event horizon

660 The left side of (29) is real only if $r \leq 2M$, namely at or inside the event
661 horizon. Whoops: *The only region on the $[r, \phi]$ slice on which we can draw*
662 *worldlines is inside the event horizon.* So what is going on *outside* the event
663 horizon? Section 3.9 answers this question; here we plot light cones on the
664 $[r, \phi]$ slice inside the event horizon. To integrate (29), use the substitution:

$$r = 2Mz^2 \quad \text{so} \quad dr = 4Mzdz \quad (30)$$

665 With this substitution, (29) becomes:

$$\frac{d\phi}{dz} = \pm \frac{4z}{(2z^2)^{1/2} (2 - 2z^2)^{1/2}} = \pm \frac{2}{(1 - z^2)^{1/2}} \quad (\text{light, } dt = 0) \quad (31)$$

666 Integrate this to obtain:

$$\phi - \phi_1 = \pm 2 \int_{z_1}^z \frac{dz}{(1 - z^2)^{1/2}} = \pm 2 [\arcsin z - \arcsin z_1] \quad (32)$$

667 Substitute back from (30) to yield the integral of (29):

$$\phi - \phi_1 = \pm 2 \left[\arcsin \left(\frac{r}{2M} \right)^{1/2} - \arcsin \left(\frac{r_1}{2M} \right)^{1/2} \right] \quad (33)$$

(light, $0 < r \leq 2M$, $0 \leq \phi < 2\pi$)

668 Light cones sprout from events at the filled dots (r_1, ϕ_1) in Figure 10.
669 Equation (33) does not give real results for $r > 2M$. However, as r approaches
670 $r_1 = 2M$ from below, the magnitude of the slopes of $d\phi/dr$ in (29) increases
671 without limit, leading to the vertical lines at $r = 2M$ in the figure.



672 **?** **Objection 15.** *Wait a minute! I thought we could draw light cones only on a*
673 *diagram with one space axis and one time axis. Figure 10 plots light cones*
674 *using two space coordinates, r and ϕ !*

3-28 Chapter 3 Curving




FIGURE 10 Light cones for different events (filled dots) on an $[r, \phi]$ slice inside and at the event horizon, showing the past (P) and future (F) of each event. Each light cone can be moved vertically, as shown. At $r = 2M$ the light moves neither inward nor outward, hence the vertical line. Because of the cyclic nature of ϕ , namely $\phi + 2\pi = \phi$, this diagram can be rolled up as a cylinder, on which the $\phi = 0$ axis and the $\phi = 2\pi$ line coincide.



675
676
677
678
679
680
681
682
683

Never assume that global coordinate separations in t , r , or ϕ tell us anything about space and time *measurements*. We favor measurement in a local inertial frame, using local coordinates—*not* global coordinates. Later we show that inside the event horizon the Schwarzschild r coordinate behaves like a time (and the Schwarzschild t coordinate behaves like a distance). So Figure 10 does describe the motion of light. The light cones in the figure fulfill one of their basic functions: For each event they divide spacetime into the past (P), the future (F), and the absolute elsewhere.

Section 3.9 Outside the event horizon: an embedding diagram on an $[r, \phi]$ slice **3-29**

3.9.4 ■ OUTSIDE THE EVENT HORIZON: AN EMBEDDING DIAGRAM ON AN $[r, \phi]$ SLICE

Freeze Schwarzschild- t ; examine stretched space.

On an $[r, \phi]$ slice:
embedding diagram
outside the
event horizon

We add a third
dimension.

Equation (29) tells us that we cannot draw light cones on the $[r, \phi]$ slice outside the event horizon. Figure 7 predicts an alternative way to visualize curved spacetime: an **embedding diagram**. Figure 12 shows the world's most famous embedding diagram, the funnel whose form we now explain and derive. Think of the $[r, \phi]$ slice outside of the event horizon as an initially horizontal rubber sheet. Here's how we create the embedding diagram: Anchor a ring at $r = 2M$ on the original flat slice, then for $r > 2M$ pull the rubber sheet upward, perpendicular to that flat surface, in such a way that the curve with $d\phi = 0$, called $Z(r)$, satisfies the equation

$$d\sigma^2 = \frac{dr^2}{\left(1 - \frac{2M}{r}\right)} \quad (\text{embedded surface profile}) \quad (34)$$

Figure 11 illustrates the resulting construction. From this figure:

$$d\sigma^2 = dZ^2 + dr^2 \quad (35)$$

From equations (34) and (35):

$$dZ^2 = d\sigma^2 - dr^2 = \frac{2M}{r - 2M} dr^2 \quad (36)$$

Take the square root of both sides of (36) and integrate the result from the lower limit at $r = 2M$:

$$Z(r) = \pm (2M)^{1/2} \int_{2M}^r \frac{dr}{(r - 2M)^{1/2}} = 2^{3/2} M^{1/2} (r - 2M)^{1/2} \quad (37)$$

Paraboloid
funnel

Spacetime only
on funnel surface

We choose the plus sign for the final expression on the right of (37) for convenience of drawing. Square both sides of (37) to obtain an equation of the form $Z^2 = Ar + B$; this shows that the funnel profile is a parabola. Rotate this curve around the vertical line $r = 0$ to create the surfaces in Figures 12 and 13. This funnel surface, with its parabola profile, is called a **paraboloid of revolution**. It is sometimes called a **gravity well** or **Flamm's paraboloid** after Ludwig Flamm, the first to identify it in 1916.

The vertical dimension in Figures 11, 12, and 13 is an artificial construct; it is *not* a dimension of spacetime. *We ourselves added this third Euclidean space dimension to help visualize Schwarzschild geometry.* Only the embedded surface represents physical spacetime where objects and people can exist. An observer posted on this paraboloidal surface is bound to stay on that surface, not because he is physically limited in any way, but because locations off the surface in these diagrams simply do not exist in physical spacetime.

The embedding diagram in Figure 13 illustrates some analytical results derived earlier in this chapter. For example:

3-30 Chapter 3 Curving




FIGURE 11 Constructing the radial profile of the funnel in Figures 12 and 13.




FIGURE 12 Space geometry visualized by distorting a slice through the center of a black hole, the result “embedded” in a three-dimensional Euclidean perspective. Adjacent circles represent adjacent shells. WE add the vertical dimension to show that the radial differential distance $d\sigma$ is greater than the differential dr (see Figure 13). Space stretching appears as a “bending” of the plane downwards into the shape of a funnel. At the throat of the funnel, where its slope is vertical, the r -coordinate is $r = 2M$.

Picturing analytical results

- 715 1. Along the radial direction, $d\sigma$ is greater than dr , as equation (35)
- 716 implies and Figure 12 illustrates.
- 717 2. The ratio $d\sigma/dr$ increases without limit as the radial coordinate
- 718 decreases toward the critical value $r = 2M$ (vertical slope of the
- 719 paraboloid at the throat of the funnel).
- 720 3. The observer constrained to the paraboloid surface cannot directly
- 721 measure the r -coordinate of any shell. He derives this r -coordinate—the
- 722 “reduced circumference”—indirectly by measuring the circumference of
- 723 the shell and dividing this circumference by 2π (Section 3.3).

Section 3.9 Outside the event horizon: an embedding diagram on an $[r, \phi]$ slice **3-31**




FIGURE 13 Projections of the embedding diagram of Figure 12. The thick curves in the side view are parabolas. WE choose the vertical coordinate for these curves in such a way that the increment along a parabola corresponds to the radial increment $d\sigma$ measured directly by the shell observer. A shell observer can exist only on the paraboloidal surface (shown edge-on as the thick curve). He can measure $d\sigma$ directly but not r or dr . He derives the r coordinate (“reduced circumference”) of a given circle by measuring its circumference and dividing by 2π . Then dr is the *computed* difference between the reduced circumferences of adjacent circles; *no* shell observer measures dr directly.

- 724 4. In contrast, the observer *can* measure the distance—call it
 725 $\sigma_{1,2}$ —between adjacent shells. He finds that this directly-measured
 726 distance is greater than the difference of their r -coordinates:
 727 $\sigma_{1,2} > r_2 - r_1$.

QUERY 2. Spacelike relation of adjacent events on an embedded surface

- A. Explain how on an embedded surface every adjacent pair of events—separated by differential global coordinates—has a spacelike relation to each other.
- B. Argue that the answer to the question, “Can a worldline (Definition 9, Section 1.5) lie on an embedding diagram?” is a resounding “NO!”
-

3-32 Chapter 3 Curving

Adjacent events on an embedding diagram have a spacelike relation.

735 In Query 1 you show that every pair of adjacent events on an embedded
736 surface has a spacelike relation to one another ($d\sigma^2 > 0$). In contrast, a stone
737 *must* move between timelike events along its worldline ($d\tau^2 > 0$). Therefore a
738 stone *cannot* move on an embedded surface. Even light—which moves along a
739 lightlike trajectory ($d\tau = 0$)—cannot move on an embedded surface. Hence an
740 embedding diagram cannot display motion at all.



741 **Objection 16.** *In a science museum I see steel balls rolling around in a*
742 *metal funnel. Is this the same as the funnel in Figure 13?*



743 No. The motion of these balls approximate Newtonian orbits provided the
744 depth at each funnel radius is proportional to the inverse of the radius,
745 which mimics the Newtonian potential energy. This is unrelated to the
746 general relativistic distortion of space near a center of gravitational
747 attraction. The cross section curve in Figure 13 is a parabola.

748 **Comment 2. Terminology: “Except on the singularity.”**

749 Neither the Schwarzschild metric, nor any other global metric we use, is valid on
750 the singularity of a black hole. On a singularity, by definition, spacetime curvature
751 increases without limit, so general relativity is not valid there. In all the global
752 coordinates we use, the non-spinning black hole has a point singularity. The
753 spinning black hole has a ring singularity in our global coordinates (Chapter 18).
754 We authors get tired of using—and you get tired of reading—the steady refrain
755 “except at the singularity.” So from now on that idea will mostly “go without
756 saying.” We will repeat the phrase occasionally, as a reminder,
757 but—please!—mentally insert the phrase “except at the singularity” into every
758 discussion of global coordinates around a black hole.



759 **Objection 17.** *So in summary, the space outside the event horizon of the*
760 *non-spinning black hole has the shape of a funnel, right? I certainly see*
761 *that funnel in textbooks and popular articles about general relativity.*



762 Here is the correct statement: “The global metric in Schwarzschild
763 coordinates leads to a funnel embedding diagram for $r > 2M$.” *Notice:*
764 This statement describes a consequence of using Schwarzschild global
765 coordinates. But it is not the consequence in *every* global coordinate
766 system. Chapter 7 introduces a global coordinate system
767 —Painlevé-Gullstrand (which we call global rain coordinates)—whose
768 global metric leads to an embedding diagram that is **flat everywhere**,
769 inside as well as outside the event horizon (Box 5, Section 7.6). The key
770 idea here is that **curvature is a property of spacetime**, not of either
771 global space coordinates alone or the global t -coordinate alone. Light
772 cone plots and embedding diagrams help us to visualize features of curved
773 spacetime, but no single diagram fully represents curved **spacetime**.
774 Sorry!




FIGURE 14 A worldtube surrounding an observer at rest in $(\phi, r/M)$ coordinates. This worldtube is bounded with slices, one of which is shaded. How “fat” the worldtube can be and still keep the the local frame of the observer inertial depends on the local spacetime curvature and the sensitivity to tides of the experiment we want to conduct.

3.10 ROOM AND WORLDTUBE

776 *Drill a hole through spacetime.*

777 We are used to the idea of experimenting or carrying out an observation in a
 778 room. A **room** is a physical enclosure, such as (1) a laboratory, (2) a powered
 779 or unpowered spaceship, or (3) an elevator with or without its supporting
 780 cables.

DEFINITION 3. Room

Definition:
room

781 A **room** is a physical enclosure of fixed spatial dimensions in which we
 782 make measurements and observations over an extended period of time.
 783

784 Thus far our room is empty; we have not yet installed the rods and clocks
 785 that allow us to record and analyze events (Figure 4, Section 5.7). However,
 786 even if the room is stationary in global r and ϕ coordinates, it changes its
 787 global t -coordinate. As it does so, the room sweeps out what we call a
 788 **worldtube** in global coordinates. Figure 14 shows the worldtube of a room at
 789 rest in r and ϕ coordinates surrounding the worldline of an observer at rest in
 790 the room.

DEFINITION 4. Worldtube

Definition:
worldtube

791 A **worldtube** is a bundle of worldlines of objects at rest in a room and
 792 worldlines of the structural components of that room. Think of a
 793 worldtube as sheathing the worldline of an observer at work in the room.
 794 Sometimes, but not always, we choose to bound the worldtube with
 795 spacetime slices, as in Figure 14.
 796

Worldtube plot
 typically curves.

797 The plot of the worldtube need not be straight, since it bounds the
 798 observer’s worldline, which typically curves in global coordinates. Figure 15
 799 shows a worldtube inside the event horizon.

3-34 Chapter 3 Curving




FIGURE 15 A worldtube inside the event horizon. The cross section of this particular worldtube is not rectangular; its sides are not slices in Schwarzschild coordinates. A horizontal or near-horizontal worldline is permitted inside the event horizon; see Figure 8.

800 In this book we prefer to make every measurement in a local inertial
 801 frame. In curved spacetime inertial frames are limited in spacetime extent.
 802 Viewed locally, each experiment takes place inside a room of limited space
 803 dimension and during a limited time lapse on clocks installed and
 804 synchronized in that room. Viewed globally, every experiment takes place
 805 within a limited segment of a worldtube.



806 **Objection 18.** *You keep saying, “In this book we prefer to make every*
 807 *measurement in a local inertial frame.” Is this necessary? Could you*
 808 *describe general relativity without using local inertial frames at all?*



809 Yes. The timelike global metric (5) delivers, on its left side, the observed
 810 wristwatch time between two events differentially close to one another. You
 811 can integrate this differential along the worldline of a stone, for example, to
 812 find the wristwatch time between two events widely separated along this
 813 worldline. A similar distant spatial separation derives from the spacelike
 814 global metric (6). All of physics hangs on events, so all of (classical,
 815 non-quantum) physics can be analyzed without local inertial frames. Our
 816 preference for measurement in local inertial frames, where special relativity
 817 rules, is a matter of taste, clarity, and convenience for us and the reader.

3.11 ■ EXERCISES**819 1. Measured Distance Between Spherical Shells**

820 A black hole has mass $M = 5$ kilometers, a little more than three times that of
 821 our Sun. Two concentric spherical shells surround this black hole. The **L**ower
 822 shell has map r -coordinate r_L ; the **H**igher shell has map r -coordinate
 823 $r_H = r_L + \Delta r$. Assume that $\Delta r = 1$ meter and consider the following four
 824 cases:

- 825 (a) $r_L = 50$ kilometers
- 826 (b) $r_L = 15$ kilometers
- 827 (c) $r_L = 10.1$ kilometers
- 828 (d) $r_L = 10.01$ kilometers
- 829 (e) $r_L = 10.001$ kilometers

- 830 A. For each case (a) through (e), use (16) to make an estimate of the
 831 radial separation σ measured directly by a shell observer. Keep three
 832 significant digits for your estimate.
- 833 B. Next, in each case (a) through (e) use the result of Sample Problem 1
 834 in Section 3.3 to find the exact distance between shells measured
 835 directly by a shell observer. Keep three significant digits for your result.
- 836 C. How do your estimates and exact results compare, to three significant
 837 digits, for each of the five cases? Give a criterion for the condition
 838 under which the estimate of part A will be a good approximation of
 839 the exact result of part B.

840 2. Grazing our Sun

841 Verify the statement in Section 3.4 concerning two spherical shells around our
 842 Sun. The lower shell, of reduced circumference $r_L = 695\,980$ kilometers, just
 843 grazes the surface of our Sun. The higher shell is of reduced circumference one
 844 kilometer greater, namely $r_H = 695\,981$ kilometers. Verify the prediction that
 845 the directly-measured distance between these shells will be 2 millimeters more
 846 than 1 kilometer. *Hint:* Use the approximation inside the front cover.
 847 (Outbursts and flares leap thousands of kilometers up from Sun's roiling
 848 surface, so this exercise is unrealistic—even if we could build these shells!)

849 3. Many Shells?

850 The President of the Black Hole Construction Company is waiting in your
 851 office when you arrive. He is waxing wroth. (“Tell Roth to wax [him] for
 852 awhile.”— Groucho Marx)

853 “You are bankrupting me!” he shouts. “We signed a contract that I would
 854 build spherical shells centered on Black Hole Alpha, the shells to be 1 meter

3-36 Chapter 3 Curving

855 apart extending down to the event horizon. But we have already constructed
 856 the total number we thought would be required and are nowhere near finished.
 857 We are running out of materials and money!"

858 "Calm down a minute." you reply. "Black Hole Alpha has an event
 859 horizon r -coordinate $r = 2M = 10$ kilometers = 10 000 meters. You agreed to
 860 build 1000 spherical shells starting at reduced circumference $r = 10\,001$
 861 meters, then $r = 10\,002$ meters, then $r = 10\,003$ meters, and so forth, ending
 862 at $r = 11\,000$ meters. So what is the problem?"

863 "I don't know. Here is our construction method: My worker robot mounts
 864 a 1-meter rod vertically (radially) from each completed shell, measures this
 865 rod in place to be sure it is exactly 1 meter long, then welds to the top end of
 866 this rod the horizontal (tangential) beam of the next spherical shell of larger
 867 r -coordinate."

868 "Ah, then your company is indeed facing a large unnecessary expense,"
 869 you conclude. "But I think I can tell you how you should construct the shells."

- 870 A. Explain to the President of the Black Hole Construction Company
 871 what his construction method should have been in order to fulfill his
 872 obligation to build 1000 correctly spaced spherical shells. Be specific,
 873 but do not be fussy.
- 874 B. Substitute the r -coordinate of the innermost shell into equation (16) to
 875 make a first estimate of the directly-measured separation between the
 876 innermost shell and the second shell, the one with the next-larger
 877 r -coordinate.
- 878 C. Using the r -coordinate of the second shell, the one just outside the
 879 innermost shell, make a second estimate of the directly-measured
 880 separation between the innermost shell and the second shell.
- 881 D. *Optional.* Use equation (18) to make an exact calculation of the
 882 directly-measured separation between the innermost shell and the one
 883 just outside it. How does the result of your exact calculation compare
 884 with the estimates of Parts B and C?
- 885 E. Determine the number of shells that the Black Hole Construction
 886 Company would have built if the President had completed the task
 887 according to his misunderstood plan.

888 4. A Dilute Black Hole

889 Most descriptions of black holes are apocalyptic; you get the impression that
 890 black holes are extremely dense objects. Of course a black hole is not dense
 891 throughout, because all matter quickly dives to the central crunch point. Still,
 892 one can speak of an artificial "average density," defined, say, by the total mass
 893 M divided by a spherical Euclidean volume of radius $r = 2M$. In terms of this
 894 definition, general relativity does not require that a black hole have a large
 895 average density. In this exercise you design a black hole with average density
 896 equal to that of the atmosphere you breathe on Earth, roughly 1 kilogram per

897 cubic meter. Carry out all calculations to one-digit accuracy—we want an
 898 estimate! *Hint:* Be careful with units, especially when dealing with both
 899 conventional and geometric units.

900 A. From the Euclidean equation for the volume of a sphere

$$V = \frac{4}{3}\pi r^3 \quad (\text{Euclid})$$

901 find an equation for the mass M of air contained in a sphere of radius
 902 r , in terms of the density ρ in kilograms/meter³. Use the conversion
 903 factor G/c^2 (Section 3.2) to express this mass in meters. (The volume
 904 formula used here is for Euclidean geometry, and we apply it to curved
 905 space geometry—so this exercise is only the first step in a more
 906 sophisticated analysis.)

907 B. Let the radius of the Euclidean spherical volume of air be equal to the
 908 map r -coordinate of the event horizon of the black hole. Assuming that
 909 our designer black hole has the density of air, what is the map r of the
 910 event horizon in terms of physical constants and air density?

911 C. Compare your answer to the radius of our solar system. The mean
 912 radius of the orbit of the (former!) planet Pluto is approximately
 913 6×10^{12} meters.

914 D. How many times the mass of our Sun is the mass of your designer
 915 black hole?

916 5. Astronaut Stretching According to Newton

917 As you dive feet first radially toward the center of a black hole, you are not
 918 physically stress-free and comfortable. True, you detect no overall accelerating
 919 “force of gravity.” But you do feel a tidal force pulling your feet and head
 920 apart and additional forces squeezing your middle from the sides like a
 921 high-quality corset. When do these tidal forces become uncomfortable? We
 922 have not yet answered this question using general relativity, but Newton is
 923 available for consultation, so let’s ask him. One-digit accuracy is plenty for
 924 numerical estimates in this exercise.

925 A. Take the derivative with respect to r of the local acceleration g in
 926 equation (13) to obtain an expression dg/dr in terms of M and r .

927 We want to find the radius r_{ouch} at which you begin to feel
 928 uncomfortable. What does “uncomfortable” mean? So that we all agree,
 929 let us say that you are uncomfortable when your head is pulled upward
 930 (relative to your middle) with a force equal to the force of gravity on
 931 Earth, $\Delta g = |g_{\text{Earth}}|$, your middle is in a local inertial frame so feels no
 932 force, and your feet are pulled downward (again, relative to your middle)
 933 with a force equal to the force of gravity on Earth $\Delta g = |g_{\text{Earth}}|$.

3-38 Chapter 3 Curving

- 934 B. How massive a black hole do you want to fall into? Suppose $M = 10$
 935 kilometers = 10 000 meters, or about seven times the mass of our Sun.
 936 Assume your head and feet are 2 meters apart. Find r_{ouch} , in meters, at
 937 which you become uncomfortable according to our criterion. Compare
 938 this radius with that of Earth's radius, namely 6.4×10^6 meters.
- 939 C. Will your discomfort increase or decrease or stay the same as you
 940 continue to fall toward the center from this radius?
- 941 D. Suppose you fall from rest at infinity. How fast are you going when you
 942 reach r_{ouch} according to Newton? Express this speed as a fraction of
 943 the speed of light.
- 944 E. Take the speed in part D to be constant from that radius to the center
 945 and find the corresponding (maximum) time in meters to travel from
 946 r_{ouch} to the center, according to Newton. This will be the maximum
 947 Newtonian time lapse during which you will be—er—uncomfortable.
- 948 F. What is the maximum time of discomfort, according to Newton,
 949 expressed in seconds?

950 *Note 1:* If you carried the symbol M for the black hole mass through these
 951 equations, you found that it canceled out in expressions for the maximum time
 952 lapse of discomfort in parts E and F. In other words, your discomfort time is
 953 the same for a black hole of *any* mass when you fall from rest at
 954 infinity—according to Newton. This equality of discomfort time for all M is
 955 also true for the general relativistic analysis.

956 *Note 2:* Suppose you drop from rest starting at a great distance from the
 957 black hole. Section 7.2 analyzes the wristwatch time lapse from any radius to
 958 the center according to general relativity. Section 7.9 examines the general
 959 relativistic “ouch time.”

960 **6. Black Hole Area Never Decreases**

961 Stephen Hawking discovered that the area of the event horizon of a black hole
 962 never decreases, when you calculate this area with the Euclidean formula
 963 $A = 4\pi r^2$. Investigate the consequences of this discovery under alternative
 964 assumptions described in parts A and B that follow.

965 **Comment 3. Increase disorder**

966 The rule that the area of a black hole's event horizon does not decrease is
 967 related in a fundamental way to the statistical law stating that the disorder (the
 968 so-called **entropy**) of an isolated physical system does not decrease. See
 969 Thorne, *Black Holes and Time Warps*, pages 422–426 and 445–446, and
 970 Wheeler, *A Journey into Gravity and Spacetime*, pages 218–222.

971 Assume that two black holes coalesce. One of the initial black holes has mass
 972 M_1 and the other has mass M_2 .

- 973 A. Assume, first, that the masses of the initial black holes simply add to
 974 give the mass of the resulting larger black hole. How does the

Section 3.11 Exercises **3-39**

975 r -coordinate of the event horizon of the final black hole relate to the
 976 r -coordinates of the event horizons of the initial black holes? How does
 977 the area of the event horizon of the final black hole relate to the areas
 978 of the event horizons of the initial black holes? Calculate the map r
 979 and area of the event horizon of the final black hole for the case where
 980 one of the initial black holes has twice the mass of the other one, that
 981 is, $M_2 = 2M_1 = 2M$; express your answers as functions of M .

982 B. Now make a different assumption about the final mass of the combined
 983 black hole. Listen to John Wheeler and Ken Ford (*Geons, Black Holes,*
 984 *and Quantum Foam*, pages 300-301) describe the coalescence of two
 985 black holes.

986 *If two balls of putty collide and stick together, the mass of*
 987 *the new, larger ball is the sum of the masses of the balls that*
 988 *collide. Not so for black holes. If two spinless, uncharged*
 989 *black holes collide and coalesce—and if they get rid of as*
 990 *much energy as they possibly can in the form of gravitational*
 991 *waves as they combine—the square of the mass of the new,*
 992 *heavier black hole is the sum of the squares of the combining*
 993 *masses. That means that a right triangle with sides scaled to*
 994 *measure the [squares of the] masses of two black holes has a*
 995 *hypotenuse that measures the [square of the] mass of the*
 996 *single black hole they form when they join. Try to picture the*
 997 *incredible tumult of two black holes locked in each other’s*
 998 *embrace, each swallowing the other, both churning space and*
 999 *time with gravitational radiation. Then marvel that the*
 1000 *simple rule of Pythagoras imposes its order on this ultimate*
 1001 *cosmic maelstrom.*

1002 Following this more realistic scenario, find the r -value of the resulting
 1003 event horizon when black holes of masses M_1 and M_2 coalesce. How
 1004 does the area of the event horizon of the final black hole relate to the
 1005 areas of the event horizons of the initial black holes?

1006 C. Do the results of both part A and part B follow Hawking’s rule that
 1007 the event horizon’s area of a black hole does not decrease?

1008 D. Assume that the mass lost in the analysis of Part B escapes as
 1009 gravitational radiation. What is the mass-equivalent of the energy of
 1010 that gravitational radiation?

1011 **7. Zeno’s Paradox**

1012 Zeno of Elea, Greece, (born about 495 BCE, died about 430 BCE) developed
 1013 several paradoxes of motion. One of these states that a body in motion
 1014 starting from Point A can reach a given final Point B only after having
 1015 traversed half the distance between Point A and Point B. But before

3-40 Chapter 3 Curving

1016 traversing this half it must cover half of that half, and so on *ad infinitum*.
 1017 Consequently the goal can never be reached.

1018 A modern reader, also named Zeno, raises a similar paradox about
 1019 crossing the event horizon. Zeno refers us to the relation between $d\sigma$ and dr
 1020 for radial separation:

$$d\sigma = \frac{dr}{\left(1 - \frac{2M}{r}\right)^{1/2}} \quad (dt = 0, d\phi = 0) \quad (38)$$

1021 Zeno then asserts, “As r approaches $2M$, the denominator on the right
 1022 hand side of (38) goes to zero, so the distance between adjacent shells becomes
 1023 infinite. Even at the speed of light, an object cannot travel an infinite distance
 1024 in a finite time. Therefore nothing can arrive at the event horizon and enter
 1025 the black hole.” Analyze and resolve this modern Zeno’s paradox using the
 1026 following argument or some other method.

1027 As often happens in relativity, the question is: Who measures what? In
 1028 order to cross the event horizon, the diving object must pass through
 1029 every shell outside the event horizon. Each shell observer measures the
 1030 incremental ruler length $d\sigma$ between his shell and the one below it. Then
 1031 the observer on that next-lower shell measures the incremental ruler
 1032 distance between that shell and the one below *it*. By adding up these
 1033 increments, we can establish a measure of the “summed ruler lengths
 1034 measured by shell observers from the shell at higher map r_H to the shell
 1035 at lower map r_L ” through which the object must move to reach the
 1036 event horizon.

1037 We integrated (38) from one shell to another in Sample Problem 1 in
 1038 Section 3.3. Let $r_L \rightarrow 2M$ in that solution, and show that the resulting
 1039 distance from r_H to r_L , the “summed ruler lengths,” is finite as
 1040 measured by the collection of collaborating shell observers. This is true
 1041 even though the right side of (38) becomes infinite exactly at $r = 2M$.

1042 Will collaborating shell observers conclude among themselves that the
 1043 in-falling stone reaches the event horizon? The present exercise shows
 1044 that the “summed ruler lengths” is finite from any shell to the event
 1045 horizon. However, motion involves not only distance but also time—and
 1046 in relativity time does not follow common expectations! What can we
 1047 say about the “summed shell time” for the passage of a diver through
 1048 the “summed shell distance” calculated above? Chapter 6, Diving, shows
 1049 that the observer on every shell measures an inertial diver to pass him
 1050 with non-zero speed, a local shell speed that continues to increase as the
 1051 diver gets closer and closer to the event horizon. Each shell observer
 1052 therefore clocks a finite (non-infinite) time for the diver to pass from his
 1053 shell to the shell below. Take the sum of these finite times—“sum”
 1054 meaning an integral similar to the integral of equation (38) carried out
 1055 in Sample Problem 1. When computed, this integral of shell times yields

1056 a finite value for the total time measured by the collection of shell
 1057 observers past whom the diver passes. Hence the group of shell observers
 1058 agree among themselves: Someone diving radially passes them all in a
 1059 finite “summed shell time” and reaches the event horizon. Thank you,
 1060 Zeno!

3.12 ■ REFERENCES

- 1062 Initial quote: John Archibald Wheeler with Kenneth Ford, *Geons, Black Holes*
 1063 *and Quantum Foam*, 1998, W. W. Norton and Company, New York, pages
 1064 296-297.
- 1065 The term *event horizon* was introduced by Wolfgang Rindler in 1956, reprinted
 1066 in in the journal *General Relativity and Gravitation*, Volume 34, Number 1,
 1067 January 2002, pages 133 through 153.
- 1068 Quotes from *The Principia* by Isaac Newton translated by I. Bernard Cohen
 1069 and Anne Whitman, University of California Press, 1999.
- 1070 References for Box 2, Section 3, “More about the Black Hole.” This box is
 1071 excerpted in part from John Archibald Wheeler, “The Lesson of the Black
 1072 Hole,” *Proceedings of the American Philosophical Society*, Volume 125,
 1073 Number 1, pages 25–37 (February 1981); J. Michell, *Philosophical*
 1074 *Transactions of the Royal Society*, London, Volume 74, pages 35–37 (1784),
 1075 cited and discussed in S. Schaffer, “John Michell and Black Holes,” *Journal*
 1076 *for the History of Astronomy*, Volume 10, pages 42–43 (1979); P.-S.
 1077 Laplace, *Exposition du système du monde*, Volume 2 (Cercle-Social, Paris,
 1078 1795), modern English translation in S. W. Hawking and G. F. R. Ellis,
 1079 *The Large Scale Structure of Space-Time*, Cambridge University Press,
 1080 Cambridge, U.K., 1973, pages 365–368; J. R. Oppenheimer and H. Snyder,
 1081 *Physical Review*, Volume 56, pages 455–459 (1939) (published the day
 1082 World War II began), quoted in Stuart L. Shapiro and Saul A. Teukolsky,
 1083 *Black Holes, White Dwarfs, and Neutron Stars: The Physics of Compact*
 1084 *Objects*, John Wiley and Sons, New York, 1983, page 338; R. P. Kerr,
 1085 *Physical Review Letters*, Volume 11, pages 237–238 (1963); E. T. Newman,
 1086 E. Couch, K. Chinnapared, A. Exton, A. Prakash, and R. Torrence, *Journal*
 1087 *of Mathematical Physics*, Volume 6, pages 918–919 (1965); S. W. Hawking
 1088 “Black Hole Explosions?” *Nature*, Volume 248, pages 30–31 (1 March 1974);
 1089 See also *Black Holes: Selected Reprints*, edited by Steven Detweiler,
 1090 American Association of Physics Teachers, New York, December 1982,
 1091 which includes reprints of papers by John Michell, Karl Schwarzschild, S.
 1092 Chandrasekhar, J. Robert Oppenheimer, and H. Snyder, Roy P. Kerr, S. W.
 1093 Hawking, and others.
- 1094 Flamm’s paraboloid, Figure 11 in Section 3.9: Ludwig Flamm (1916).
 1095 ”Beiträge zur Einstein’schen Gravitationstheorie”. *Physikalische Zeitschrift*
 1096 Volume 17, pages 448-454.

3-42 Chapter 3 Curving

1097 Some items in Box 5 adapted from Misner, Thorne, and Wheeler,
1098 GRAVITATION, W. H. Freeman Company, 1970, San Francisco (now New
1099 York), page 671
1100 Quote *Black holes just didn't "smell right"* from Kip Thorne, *Black Holes and*
1101 *Time Warps: Einstein's Outrageous Legacy*, New York, W. W. Norton,
1102 1994, pages 134 and 137.

1103 Download file name: Ch03Curving170511v1.pdf